

# **Wetland Ecology: Examining the Role of Wetlands in Water Quality Improvement and Biodiversity Conservation**

<sup>1</sup> Dr. Jo Rivero, Universidad Científica Del Sur, Peru.

**Abstract:** Because of agribusiness, supplement improvement has a significant impact on the water quality in many stream catchments and river bowls. Since there is evidence at the site level that riparian zones and floodplain wetlands are successful at removing supplements from through-streaming water, water-asset directors generally are considering the predicted role of these wetlands in further improving stream-water quality. Water quality improvement has been shown to occur at the landscape level where groups of aquatic and wetland ecosystems are hydrologically connected and at the level of individual systems. At the landscape level, one example of the influence that groups of wetlands and shallow aquatic ecosystems have on water quality comes from research on beaver have shown that beaver influence water quality over large areas of boreal landscapes and that the impacts are dynamic and long lasting. Research on beavers has demonstrated that they have a dynamic and enduring impact on water quality across vast swaths of boreal landscapes. This is one example of how groupings of wetlands and shallow aquatic ecosystems affect water quality at the landscape level. Beavers modify hydrologic patterns and produce wetlands and shallow impoundments that are perfect for changing the quality of water, especially by affecting the dynamics of carbon and nitrogen. Wetlands are essential for regulating climate change and promoting biodiversity protection. In the past, wetlands were seen as wastelands, which led to their degradation in a number of locations. Therefore, the preservation of the wetland ecosystem necessitates strict environmental regulations.

**Keywords:** Wetland Ecosystem; Water Quality Improvement; Biodiversity Conservation.

(Submitted: March 17, 2024; Revised: April 09, 2024; Accepted: May 13, 2024; Published: June 28, 2024)

## **I. Introduction**

Wetlands have characteristics of both aquatic and terrestrial ecosystems and form where they meet. Although they may differ greatly in terms of appearance and species makeup, water inundation is a common trait that is mirrored in soil processes and biota adaptations. As a result, wetlands can be found anywhere there is water, from the interior of the mainland to sour coastal areas, but they are typically connected to freshwater. When water immersion produces soils that are overpowered by anaerobic processes, a climate known as a wetland is created. As a result, the biota—especially established plants—adapts to flooding. Subarctic peatlands and tropical mangrove swamps are both included in this broad description. There is a cause (water flooding), a proximate effect (soil oxygen levels being lowered), and a secondary effect (the biota having to withstand both the direct impacts of flooding and the secondary effects of anaerobic conditions) in just one sentence of definition. While it may not be the perfect definition, it is the only one that will get us started. We will begin with this straightforward concept because numerous scientists, attorneys, agencies, and groups have tried to define wetlands. Since wetlands need water, the distribution of water on Earth is a clear place to start. Fresh water makes up a far lesser portion. A distillery is powered by solar heat, which extracts water vapor from the oceans and returns it to the land as precipitation. Certain types of wetlands, such as salt marshes in higher latitudes and mangrove swamps in equatorial regions, develop near the ocean's shore. However, freshwater ecosystems make up the majority of wetlands. They happen in places where rainwater collects before returning to the ocean. You will frequently come across documents that refer to "coastal wetlands" and "interior wetlands," and some people consider the difference between freshwater and saltwater wetlands to be crucial. Although salinity undoubtedly has a significant role in determining the types of plants and animals that exist, we will try our best in this book to consider wetlands as a single ecosystem Figure 1. The majority of life, including freshwater life, has a chemical composition that is closer to that of the ocean than freshwater. However, it seems that the majority of life that exists in fresh water today did not begin there; rather, it first adapted to land before moving on to fresh water. Naturally, fish were an exception.



Figure 1: Wetlands

One of nature's traits and a unique feature of living systems is biodiversity. According to the Show on Natural Variety, biodiversity is "the inconstancy among living life forms from all sources including, bury alia, earthbound, marine, and other amphibian biological systems and the environmental edifices of which they are part, including variety inside species, among species, and of biological systems." This definition is the most well-known definition of biodiversity. Various definitions have been put out to highlight the variation at various levels of association, ranging from settings to attributes. In any event, due to the global interest in the social, economic, political, legitimate, and other aspects of biodiversity and its destruction, the word has been used pretty widely to obligate various vested parties. Although interest in many facets of biodiversity has skyrocketed in recent years, the majority of research has concentrated on terrestrial or marine habitats. Only recently has attention begun to be paid to the biodiversity found in inland aquatic habitats. Wetlands have been well studied for their ecology, management, conservation, and restoration among inland aquatic ecosystems; nevertheless, until recently, the concerns pertaining to their biodiversity were mostly ignored.

## II. Literature Review

Wetlands serve significant economic activities like tourism and offer special and irreplaceable functions connected to a variety of ecosystem services vital for biodiversity conservation, mitigating the effects of climate change, and promoting human well-being (Durigan et al., 2022). Wetlands are impacted by human activity both directly and indirectly. Dredging, filling, and draining wetlands on purpose, including through the state and federal wetlands permitting process, are examples of direct impacts. Activities that change the hydrological system, increase pollution loadings, or induce buffer encroachment due to urbanization and development are examples of indirect consequences. With a strict water quality monitoring program, these effects can be reduced and tracked (Ballut-Dajud et al., 2022). Wetlands can be severely impacted by poor water quality. The data gathered from water quality evaluations can be utilized to comprehend the impact of human activities on wetlands. The variety and abundance of the plants and animals are influenced by the

pH, mineral concentration, and nutrient content. Each nutrient has a distinct purpose in the ecosystem of wetlands and, if concentrations are not routinely checked, can lead to pollution through eutrophication. Small amounts of nutrients, trace metals, and other substances can be taken up by certain wetland plants from the soil water and incorporated into their own tissue, which can then be recycled in the wetland through decomposition (Brisson et al., 2020). The Water Quality Index (WQI) measures eight elements to assess the quality of the water. Temperature, pH, dissolved oxygen (DO), biochemical oxygen demand (BOD), turbidity, total phosphorus, nitrates, and *E. coli* are some of these variables. Wetlands are difficult to classify due to the existing lack of agreement on what constitutes a wetland, and various systems have been devised. The Ramsar Characterisation Framework for Wetland Type is the one that is used the most widely on a general basis. It divides wetlands into three categories: inland, marine/waterfront, and man-made, based on their area, vegetation type, hydrological conditions, and regular or man-made climate. These three fundamental Ramsar classifications are further separated into 42 distinct types of wetlands based on a number of factors, such as substrate and water type. Another well-known grouping strategy is a multi-leveled characterisation framework based on hydrological, geomorphological, synthetic, and organic components (Hammer & Bastian, 2020).

### III. Impact of Wetlands in Improving Water Quality

Wetlands serve as nature-based arrangements (NBS) to reduce the levels of foreign substances in surface streams by mitigating the detrimental effects of soil disintegration, overflow, and wastewater tainting on water quality. Wetlands prevent the deterioration of water quality by capturing pollutants from mechanical processes (such as sedimentation and filtration), adsorption on the substrate, biosorption, and other complex and interrelated interactions between plants and microorganisms, as well as sanitisation from UV radiation from daylight. They may remove a variety of natural and inorganic substances from contaminated water, such as vitamins, heavy metals, pesticides, hydrocarbons, xenobiotics, and antimicrobials (Moshiri, 2020). These pollutants can generally be found in the most temperate water in rural areas, urban surfaces, civic wastewater, landfill leachate, hydroponics effluent, and some contemporary wastewaters. Because there is a limit to how much can be added before the normal synthetic and plant processes get overworked and debased, normal wetlands have a somewhat limited ability to handle increasing contaminants. Furthermore, wetlands can be used to recycle wastewater and water for other uses (such as water systems), which is especially important in areas with limited water supplies (Scholz, 2023). In the last ten years, microbial power devices have been combined with artificial wetlands to create a novel wastewater treatment technique that produces bioenergy.

### IV. Wetland Ecology: Conservation and Management

In the past, temperate and tropical regions received distinct treatment for wetlands and their biota. In contrast to the West, where they were frequently derided for harboring disease and ghosts, wetlands were revered for their significance in socioeconomic and cultural life. Understanding traditional management techniques and the intricate relationships between humans and wetland biota are essential for the protection and management of wetlands (Irvine et al., 2022). Activities, as well as a straightforward understanding of the environmental mechanisms that sustain them. Wetland management objectives must be well-defined and informed by the specific wetland's natural characteristics. In general, a wetland's several functions are incompatible with one another. For instance, some wetlands aid in the treatment of wastewater that passes through them, but excessive discharge can harm biodiversity by releasing harmful compounds or nitrogen enrichment, making the wetland unsuitable for enjoyment and water supply. Business fishing and hydroponics frequently clash with the interests of traditional fisheries and other biodiversity. Wetlands have been the focus of protection efforts thus far, typically based on ideas and methods developed for terrestrial settings without determining the primary distinction between the two. Wetlands' form and potential are determined by the medium, or water, just like in other freshwater biological systems; biota and other elements play an optional role. A variety of environmental factors typically keep wetlands in their younger phases of succession, away from the climax terrestrial condition (Yu et al., 2020). The majority of wetland

biota sustain a particular measure of annual water level fluctuation, comparable to that measured in floodplains and lake littoral zones. There are long periods of drought and flooding in different types of wetlands. Anthropogenic alterations to hydrology could lead to a significant loss of biodiversity. Protecting nearby high country and large aquatic environments, as well as maintaining a high degree of living space variety within the wetland, are also essential for preserving biodiversity in wetlands shown in Figure 2.




Figure 2: Functions of Wetland Ecosystem

## V. Conclusion

A vast assortment of plant and creature species can be viewed as in wetlands, making them hotspots for biodiversity. They offer migratory fish, amphibians, birds, and many other wildlife species a place to live. Wetlands' varied vegetation reduces the effects of climate change by sequestering carbon. Wetlands must be preserved in order to protect biodiversity, stop the extinction of species, and preserve ecological balance. Wetlands provide substantial economic advantages in addition to their ecological significance. They give local populations access to sources of income by promoting the tourism, agricultural, and fishing sectors. Wetlands support the sustainability of natural resources, agricultural output, and soil fertility. Long-term financial benefits and environmental resilience can result from the preservation and restoration of wetlands. Wetlands are undermined by various variables, like contamination, climate change, and habitat loss, notwithstanding their significance. Wetland loss and degradation are frequently caused by infrastructure development, urbanization, and agricultural growth. Wetland ecosystems are further threatened by pollution from waste dumping, agricultural chemicals, and industrial runoff. These issues are made worse by climate change, as wetland health is impacted by changing rainfall patterns and rising temperatures.

## References

- [1] Durigan, G., Munhoz, C. B., Zakia, M. J. B., Oliveira, R. S., Pilon, N. A., do Valle, R. S. T., ... & Pott, A. (2022). Cerrado wetlands: multiple ecosystems deserving legal protection as a unique and irreplaceable treasure. *Perspectives in ecology and conservation*, 20(3), 185-196. <https://doi.org/10.1016/j.pecon.2022.06.002>

- [2] Ballut-Dajud, G. A., Sandoval Herazo, L. C., Fernández-Lambert, G., Marín-Muñiz, J. L., López Méndez, M. C., & Betanzo-Torres, E. A. (2022). Factors affecting wetland loss: A review. *Land*, 11(3), 434. <https://doi.org/10.3390/land11030434>
- [3] Brisson, J., Rodriguez, M., Martin, C. A., & Proulx, R. (2020). Plant diversity effect on water quality in wetlands: a meta-analysis based on experimental systems. *Ecological Applications*, 30(4), e02074. <https://doi.org/10.1002/eap.2074>
- [4] Hammer, D. A., & Bastian, R. K. (2020). Wetlands ecosystems: natural water purifiers?. In *Constructed wetlands for wastewater treatment* (pp. 5-19). CRC Press.
- [5] Moshiri, G. A. (2020). *Constructed wetlands for water quality improvement*. CRC Press.
- [6] Scholz, M. (2023). *Wetlands for water pollution control*. Elsevier.
- [7] Irvine, K., Dickens, C., Castello, L., Bredin, I., & Finlayson, C. M. (2022). Vegetated wetlands: from ecology to conservation management. In *Fundamentals of Tropical Freshwater Wetlands* (pp. 589-639). Elsevier. <https://doi.org/10.1016/B978-0-12-822362-8.00023-2>
- [8] Yu, W., Ji, R., Han, X., Chen, L., Feng, R., Wu, J., & Zhang, Y. (2020). Evaluation of the biodiversity conservation function in Liaohé Delta wetland, northeastern China. *Journal of Meteorological Research*, 34(4), 798-805. <https://doi.org/10.1007/s13351-020-9186-7>