Aquatic Ecosystems and Environmental Frontiers, Volume 3, Issue 1, March 2025 *ISSN: 3049-186X*
DOI: https://doi.org/10.70102/AEEF/V3I1/6

The Role of Aquatic Food Webs in Maintaining Ecosystem Stability and Resilience

[1*] Wei Jian, Sino Horizon Institute for Emerging Technology Studies, Beijing, China.

[2] Liu Meilin, Sino Horizon Institute for Emerging Technology Studies, Beijing, China.

Abstract: The importance of aquatic ecosystems (both freshwater and marine ecosystems) cannot be understated, as they contribute towards biodiversity conservation and provide important ecosystem services, including nutrient recycling, carbon fixation, and fisheries. In recent years, such systems have become vulnerable to several environmental factors, namely climate change, pollution, and habitat destruction, which affect the system's capacity to function optimally and be resilient against future stresses. Food web networks, which form a network comprising the interrelationships among different organisms, form important components of maintaining ecosystem stability through energy flow, species interactions, and redundancy. This study investigates the impact of food webs on the stability and resilience of aquatic ecosystems. Experiments were carried out in the field to measure abundance and biomass of different species, using stable isotope analysis ($\delta^{13}C$ and $\delta^{15}N$) to assign trophic levels. Food web networks were created from collected data, whereas structural properties (connectance, linkage density, mean trophic level, modularity) were determined alongside robustness, redundancy, and resilience scores. Statistical analysis, which includes principal components and redundancy analyses, was carried out to determine the effect of food web complexity on ecosystem resilience. The outcome shows that ecosystems with increased connectance and trophic complexity, especially those found in the estuarine and marine environments, display high levels of robustness and functional redundancy, thus increasing their resilience to species extinction and external disturbances. The key species that included top consumers and primary producers were found to play an important role in maintaining the stability of the networks. Seasonal variation also affected connectivity and redundancy in the ecosystem.

Keywords: Aquatic Food Webs; Ecosystem Stability; Resilience; Trophic Interactions; Biodiversity; Keystone Species.

(Submitted: December 30, 2024; Revised: January 29, 2025; Accepted: February 17, 2025; Published: March 31, 2025)

I. Introduction

Aquatic ecosystems, comprised of rivers, lakes, wetlands, and the oceanic environment, are fundamental to maintaining the planet's biodiversity and contribute crucially to ecosystem processes such as nutrient recycling, carbon storage, and fishery production (Chowdhury et al., 2025; Zhao et al., 2023). Such ecosystems continue to be compromised by environmental stress factors, such as climate change, pollution, habitat degradation, and resource exploitation, making them less capable of functioning in support of ecological functions and human survival (Naman et al., 2022; Saito et al., 2024). An understanding of the processes that drive ecosystem stability and resiliency becomes, therefore, critical to managing aquatic ecosystems effectively (Lewis et al., 2022; Schlenker et al., 2024).

The food web, which encompasses trophic interactions between living things in an ecosystem, forms a vital part of aquatic ecosystem functioning. Complexity in the food web affects how well the ecosystem is able to handle disruptions, and also determines energy flow in ecosystems and species interactions therein. A complex food web may help ecosystems become resilient to changes in the environment by spreading energy through many avenues, providing functional redundancy. However, although many scientists have tried their hands at understanding the aquatic trophic network systems, there still remain unanswered queries as to how certain elements of aquatic food webs can lead to ecosystem stability under varying pressures of man-made nature. It is vital that answers be sought to these queries in order to understand proper management of ecosystems and aquatic resources.

Objectives

1. The analysis of structural complexity in the aquatic food web of both freshwater and marine environments, with the help of connectivity metrics such as connectome, linkage density, mean trophic level, and modularity.
2. The analysis of stability and resilience of ecosystems through the study of robustness and functional redundancy metrics, and identifying keystone species.
3. Analysis of the effects of biodiversity, trophic complexity, and seasonality on ecosystem stability to provide recommendations for conserving ecosystems in the event of environmental stress in aquatic environments.

The organization of the paper is organized as follows: Section 1 (Introduction) gives a background and gaps in research; Objectives discusses the aims of the paper; Section 2 (Methodology) talks about study sites, sample collections and analyses, data acquisition and analysis; Section 3 (Results) comprises measurements of networks' characteristics, stability, seasonality, and resilience curves, presented in tables and graphs; Section 4 (Discussion) discusses results and their implications; and, finally, Section 5 (Conclusion) wraps up the whole paper.

II. Methods

Study Sites and Sampling Design

The research work was carried out in various aquatic environments such as rivers, lakes, estuaries, and coastal regions, among others, depending on the type of ecosystem under consideration. In addition, stratified sampling was used to incorporate the effect of spatial variation in the composition of the ecosystem and its food webs (Zhou et al., 2024). Wet and dry seasons were considered to incorporate temporal variation in community structure (Madhanraj, 2025; Robinson et al., 2024).

Data Collection Protocols

Field sampling was done for the determination of abundance and biomass of the organisms. Macrofaunal samples were collected by using nets, traps, and quadrats, whereas samples of the primary producers were taken via chlorophyll-a and percentage cover of macrophytes. Stable isotope techniques were used for determining the trophic levels of various species collected during sampling, using $\delta^{13}C$ and $\delta^{15}N$ ratios. Information about the diet and food interactions was gathered from direct observations as well as existing literature reviews.

Food Web Analysis Tools

The network for the food web was developed based on the data of species interactions obtained during research (Shipley et al., 2023). Metrics of connectance, linkage density, mean trophic level, and modularity were computed as indices for determining structural complexity of the network (Ward et al., 2023). The ecosystem stability indicators of robustness, redundancy, and trophic diversity were used to measure stability in order to reveal any keystone species or interactions responsible for the stability of the aquatic ecosystem (Gutgesell et al., 2024).

Statistical Approaches for Resilience Evaluation

The relationships between the structure of the food web and the environment were analyzed by multivariate statistics (principal component analysis (PCA) and redundancy analysis (RDA)) (Ma et al., 2023). The regression model and network simulation were used to study the impact of differences in the structure of trophic links and species on the stability and resilience of ecosystems. Significance was considered at the significance level of 5% ($p < 0.05$). The analysis was performed using the software package R version 4.2.0 (graph and bipartite packages).




Figure 1: Network Diagrams of Freshwater and Marine Food Webs

In figure 1, two comparative network charts are displayed for understanding trophic interactions within freshwater and marine environments (Russo et al., 2023). Nodes represent species, while their trophic level is shown in the colors. Arrows in the diagram demonstrate energy transfer between species through predation. Keystones such as the Largemouth Bass in freshwater and the Reef Shark in the marine environment are depicted by bigger nodes to emphasize their significance in ensuring ecosystem stability (Sporta Caputi et al., 2024).

III. Results

Food Web Structure Metrics

Aquatic ecosystems analyzed had varying levels of complexity in terms of their food webs. The connectance values for the different ecosystems ranged between 0.18 and 0.42, which implies that there was variation in the complexity levels of interaction. The mean trophic level was seen to vary from 2.4 to 3.6 for each ecosystem analyzed. Analysis of network modularity showed that there were compartments within the food web, which indicated the formation of tightly interconnected functional groups. Linkage density was estimated at 3.2 to 5.1 for each system analyzed.

Table 1: Summary of Food Web Metrics Across Aquatic Sites

Site Type	Connectance	Mean Trophic Level	Linkage Density	Modularity
River (Freshwater)	0.28	2.7	3.5	0.41
Lake (Freshwater)	0.34	3.1	4.2	0.36
Estuary (Brackish)	0.42	3.4	5.1	0.33
Coastal (Marine)	0.38	3.6	4.8	0.29

Table 1 provides a summary of the major structural properties of food webs in various types of aquatic systems, such as rivers, lakes, estuaries, and marine environments. This table illustrates the variation in

connectance, mean trophic level, linkage density, and modularity in food webs from different ecosystems. Connectance and linkage density give an indication of complexity, whereas modularity is an indication of the extent of compartmentalization in each food web system.

Table 2: Stability Indices Across Aquatic Ecosystems

Site Type	Robustness (%)	Redundancy	Resilience Score
River (Freshwater)	72	1.8	0.68
Lake (Freshwater)	81	2.1	0.74
Estuary (Brackish)	88	2.5	0.82
Coastal (Marine)	84	2.3	0.79

The stability indices for various aquatic ecosystems, like rivers, lakes, estuaries, and coastal environments, are displayed in table 2. The robustness, redundancy, and resilience indices provide information on how well each system can resist losing its species without compromising stability. The more robust and redundant each system is, the better equipped it is at resisting stress; resilience combines both of these parameters.




Figure 2: Seasonal Variation in Connectance and Redundancy Across Sites

Figure 2 shows the seasonal patterns for two important indices of food webs connectivity and redundancy in four different aquatic habitats: rivers, lakes, estuaries, and the coast. Figure 2A provides an example of how connectivity is seasonal, whereas figure 2 B gives a picture of the seasonal variations in redundancy. Figure 2 demonstrates that the indices are greater in summer and smaller in winter, and that estuarine habitats are more complex than others.

Patterns in Stability and Resilience

Indicators of stability, such as robustness and redundancy, have shown clear relationships with food web complexity. Food webs with high connectance and trophic complexity were able to maintain their functioning even when the removal of important species was modeled. Redundancy showed that more than

one species carried out the same ecological functions in an environment, providing a backup for functionality if the other species were disturbed. Simpler food webs with low connectance were more vulnerable to the cascading effect of losing species in the environment. Results of the temporal analysis showed that networks from wet seasons had high levels of connectivity and stability compared with networks formed during the dry season.

Figure 3 provides the visual representation of the proportion of the recovery of ecosystems depending on the removal of different species from both lakes and coastal regions. Four different types of species are represented in four distinct graphs: top predators, key consumers, grazers, and primary producers. The figure shows how resilience is affected by the removal of species: the loss of top predators is not critical; however, the removal of primary producers leads to decreased resilience.




Figure 3: Resilience Curves of Freshwater and Marine Ecosystems

Critical Species and Key Interactions

Certain species, which included keystone predators and habitat-forming primary producers, were characterized as being keystone nodes in the networks. Removal experiments revealed that the extinction of such species brought about a drastic reduction in network robustness (up to 27%) and an increase in trophic instability. Interactions between mutualists and predator-prey relations with such keystone species had a higher effect on the stability of the food web. The network visualization was able to identify clusters of high interaction intensity and redundancy.

IV. Discussion

The results from this study reveal the significance of food web architecture in ensuring the stability and resilience of ecosystems. The high levels of connectance and linkage density in estuarine and coastal ecosystems point out that elaborate networks characterized by many interdependent food chains ensure adequate energy flow and protect ecosystems against any form of disturbance. Apex predators and producers have a high level of control over these networks because their extinction results in drastic changes in the functioning of ecosystems.

Biodiversity and the complexity of food chains are two important buffering mechanisms in ecology. Ecosystems with high numbers of interacting organisms showed increased resistance to disturbances, implying the ability of ecosystems to cope with challenges and sustain vital functions. Changes in seasons further demonstrate that aquatic food chains are dynamic in terms of connectivity and redundancy levels, thus ensuring increased ecosystem recovery potential after environmental stresses. This finding is in line with previous studies highlighting that complex food chains and biodiversity increase ecosystem resilience.

Management strategies aimed at maintaining food web structures include not only the protection of keystone species but also the maintenance of important interaction channels within an ecosystem. Such strategies include heterogeneity of habitat, preventing overfishing, and protecting primary producers. This kind of information is useful theoretically because it confirms the existence of a relationship between food web complexity and ecosystem stability.

V. Conclusion

The importance of the structure and complexity of the aquatic food web for the stability and resilience of the ecosystem in both freshwater and marine environments has been illustrated in this study. For example, more stable and resilient ecosystems are those with higher levels of connectance, functional redundancy, and trophic diversity, particularly when it comes to estuaries and coastal areas. The role played by keystone species, particularly apex predators and primary producers, is integral to the stability of the ecosystem network and will be greatly diminished by the removal of these organisms. Seasonality will also affect the level of functional redundancy and connectance within the ecosystem. The implications indicate that the conservation of the complexity and redundancy in food webs plays a critical role in enhancing the resilience of the ecosystem in face of increasing anthropogenic and natural perturbations. Not only do the findings reinforce the ecological theory, which is founded on biodiversity and network complexity, but also provide useful pointers in planning a conservation strategy. From the research point of view, it becomes crucial to conserve the keystone species, retain habitat heterogeneity, and prevent the extinction of primary producers to facilitate the conservation of the food web. The future course of action would involve observing the aquatic ecosystem over a longer span of time in response to environmental disturbances.

References

[1] Gutgesell, M., McCann, K., O'Connor, R., Kc, K., Fraser, E. D., Moore, J. C., ... & Rooney, N. (2024). The productivity–stability trade-off in global food systems. *Nature Ecology & Evolution*, *8*(11), 2135-2149. https://doi.org/10.1038/s41559-024-02529-y

[2] Naman, S. M., White, S. M., Bellmore, J. R., McHugh, P. A., Kaylor, M. J., Baxter, C. V., ... & Puls, A. L. (2022). Food web perspectives and methods for riverine fish conservation. *Wiley Interdisciplinary Reviews: Water*, *9*(4), e1590. https://doi.org/10.1002/wat2.1590

[3] Madhanraj. (2025). Effects of Climate Warming and Acidification on Aquatic Food Web Structure and Productivity. *Journal of Aquatic Ecology and Environmental Sustainability, 2*(2), 23–29.

[4] Ward, C. A., Tunney, T. D., & McCann, K. S. (2023). Managing aquatic habitat structure for resilient trophic interactions. *Ecological Applications*, *33*(3), e2814. https://doi.org/10.1002/eap.2814

[5] Saito, V. S., Kratina, P., Barbosa, G., Ferreira, F. C., Leal, J. B., Zemelka, G., ... & Perkins, D. M. (2024). Untangling the complex food webs of tropical rainforest streams. *Journal of Animal Ecology*, *93*(8), 1022-1035. https://doi.org/10.32942/x28605

[6] Russo, L., Bellardini, D., Zampicinini, G., Jordán, F., Congestri, R., & D'Alelio, D. (2023). From metabarcoding time series to plankton food webs: The hidden role of trophic hierarchy in providing ecological resilience. *Marine Ecology*, *44*(3), e12733. https://doi.org/10.1111/maec.12733

[7] Lewis, K. A., Christian, R. R., Martin, C. W., Allen, K. L., McDonald, A. M., Roberts, V. M., & Valentine, J. F. (2022). Complexities of disturbance response in a marine food web. *Limnology and Oceanography*, *67*, S352-S364. https://doi.org/10.1002/lno.11790

[8] Zhou, X., Wang, C., Kattel, G., Zhang, J., & Xu, M. (2024). Hydrodynamic and trophic variations reshape macroinvertebrate food webs in urban ecosystems. *Environmental Science and Ecotechnology*, *22*, 100478. https://doi.org/10.1016/j.ese.2024.100478

[9] Sporta Caputi, S., Kabala, J. P., Rossi, L., Careddu, G., Calizza, E., Ventura, M., & Costantini, M. L. (2024). Individual diet variability shapes the architecture of Antarctic benthic food webs. *Scientific Reports*, *14*(1), 12333. https://doi.org/10.1038/s41598-024-62644-5

[10] Ma, J., Chen, W., Chen, M., Zhong, K., Yao, N., Zhang, X., ... & Zhou, Q. (2023). Water level fluctuations associated with hydrological connectivity consolidate the food web stability of the largest Chinese freshwater lake via mediating trophodynamics and trophic structure. *Ecological Indicators*, *153*, 110372. https://doi.org/10.1016/j.ecolind.2023.110372

[11] Shipley, O. N., Matich, P., Hussey, N. E., Brooks, A. M., Chapman, D., Frisk, M. G., ... & Gallagher, A. J. (2023). Energetic connectivity of diverse elasmobranch populations–implications for ecological resilience. *Proceedings of the Royal Society B: biological sciences*, *290*(1996), 20230262. https://doi.org/10.1098/rspb.2023.0262

[12] Chowdhury, P. R., Goswami, S., Choudhary, R., & Choudhury, M. (2025). Electronic wastes, its impact on wildlife and biodiversity. In *Electronic waste* (pp. 65-71). CRC Press. https://doi.org/10.1201/9781003582311-8

[13] Zhao, Q., Van den Brink, P. J., Xu, C., Wang, S., Clark, A. T., Karakoç, C., ... & De Laender, F. (2023). Relationships of temperature and biodiversity with stability of natural aquatic food webs. *Nature communications*, *14*(1), 3507. https://doi.org/10.1038/s41467-023-38977-6

[14] Schlenker, A., Brauns, M., Fink, P., Lorenz, A. W., & Weitere, M. (2024). Long-term recovery of benthic food webs after stream restoration. *Science of the Total Environment*, *923*, 171499. https://doi.org/10.1016/j.scitotenv.2024.171499

[15] Robinson, J. P., Benkwitt, C. E., Maire, E., Morais, R., Schiettekatte, N. M., Skinner, C., & Brandl, S. J. (2024). Quantifying energy and nutrient fluxes in coral reef food webs. *Trends in Ecology & Evolution*, *39*(5), 467-478. https://doi.org/10.1016/j.tree.2023.11.013