Aquatic Ecosystems and Environmental Frontiers, Volume 3, Issue 1, March 2025 *ISSN: 3049-186X*
DOI: https://doi.org/10.70102/AEEF/V3I1/7

Biotic and Abiotic Interactions in Freshwater Ecosystems and their Role in Ecosystem Functioning

[1*] Ahmad Fauzi, Malaysian Institute for Smart Technology Research, Kuala Lumpur, Malaysia.

[2] Nur Aisyah, Malaysian Institute for Smart Technology Research, Kuala Lumpur, Malaysia.

Abstract: Freshwater biomes such as lakes, wetlands, and rivers are important for reasons such as species richness, nutrient turnover, and regulation of water purity. Relationships among organisms in an ecosystem, including producers, consumers, and decomposers, as well as abiotic components such as temperature, hydrology, oxygen, and nutrient content, influence the nature of ecosystems. In this literature review, an analysis of current research on the interaction of biotic and abiotic factors and their impacts on ecosystems is conducted. Research articles were selected according to their relevance, inclusion of both biotic and abiotic factors, and evaluation of ecosystem functioning within fifteen years ago to date. It was found that changes in abiotic factors affected species and trophic relationships in the community, and in turn, interactions between organisms regulated ecosystem-level processes. It was concluded that nutrient loading, variability in flow, and temperature were important abiotic regulators, and high functional diversity and complexity of food webs contributed to greater ecosystem stability and resilience. The integration of these observations underscores the relevance of taking into account the role of bottom-up and top-down processes in the management of freshwater ecosystems. Some of the implications of these observations for the conservation and management of freshwater ecosystems include the maintenance of the natural hydrology of the systems, the preservation of water quality, the maintenance of ecological heterogeneity, and the restoration of ecological functions.

Keywords: Freshwater Ecosystems; Biotic–Abiotic Interactions; Nutrient Cycling; Ecosystem Resilience; Species Diversity; Productivity; Conservation Management.

(Submitted: December 31, 2024; Revised: January 30, 2025; Accepted: February 18, 2025; Published: March 31, 2025)

I. Introduction

Freshwater ecosystems, comprising rivers, lakes, and wetlands, are important parts of the biosphere that contribute to essential ecosystem services such as providing habitat for biodiversity, cleaning water, and recycling nutrients (Atkinson et al., 2017). Freshwater ecosystems are crucial for sustaining complex communities by regulating various hydrologic and biogeochemical cycles (Setubal et al., 2020).

The biotic aspects of the freshwater environment include all living things, such as primary producers, microorganisms, invertebrates, and fish, and their interactions, such as predation, competition, and mutualism, define the nature of their community and its function (Rodrigues-Filho et al., 2023). The interactions between different trophic levels and biodiversity define how ecosystems work and recycle their nutrients (Manthila et al., 2025).

Abiotic aspects include all physical and chemical conditions, such as temperature, water flow, availability of oxygen, pH level, and nutrients. All the aforementioned abiotic characteristics create an environment that defines the distribution and functioning of living things, as well as affects the interaction among species (Birk et al., 2020).

Knowledge on how biotic and abiotic factors interact is important in gaining knowledge on how freshwater systems work. The interaction of the two influences processes such as primary production, nutrient recycling, and ecosystem stability, as well as the response of the system to environmental stressors such as climate change and human pollution (Vaughn, 2010). It is therefore very important to understand both the biotic and abiotic aspects of an ecosystem (Nõges et al., 2016).

Objectives

1. Evaluate the impact of abiotic components on the community structures and functions of aquatic biotic communities.
2. Determine the effects of interaction between biotic and abiotic components on various ecological processes like productivity, recycling of nutrients, etc.
3. Present ideas for the conservation and restoration of stressed freshwater ecosystems based on evaluation.

Structure of the article: Section 1 (Introduction) – introduction to freshwater ecosystems and their biotic and abiotic components; Section 2 (Methods) – description of the methodology used in this study; Section 3 (Results) – results on abiotic patterns, biotic distribution, and interaction effects; Section 4 (Discussion) – interpretation of the results taking into account ecological processes and resiliency; and finally, Section 5 (Conclusion).

II. Methods

The objective of this literature review is to systematically synthesize the existing information on the interaction between biotic and abiotic entities within fresh water and its effect on the ecosystem's functionality. Articles were selected depending on their relevance to freshwater ecosystems, presence of biotic and abiotic elements, and focus on ecosystem functions such as productivity and resilience.

Figure 1 The above diagram highlights the relationship between abiotic elements (water availability, chemistry, temperature, light, weather conditions, and land structure), biotic elements (producers, consumers, decomposers, and food chains), and processes resulting from ecosystem functioning (primary productivity, nutrient cycling, energy flow, water quality regulation, and ecosystem stability). The arrows represent causal connections and feedback loops, emphasizing how abiotic elements affect the makeup of the community and processes occurring in ecosystems, while at the same time, the activities of the biotic components influence the abiotic environment.




Figure 1: Conceptual Model of Biotic–Abiotic Interactions in Freshwater Ecosystems

The review concentrated on river, lake, and wetland ecosystems in different geographical areas in order to take into account various hydrological and climatic conditions. The biotic variables that were analyzed were species diversity, community structure, trophic relationships, and functional groups. Hydrological variables such as temperature, hydrology, oxygen concentration, nutrient content, pH value, and light availability have been identified as abiotic variables.

Studies that were included in the review had to satisfy specific selection criteria. These criteria were that the paper was peer-reviewed, had to be published no earlier than 15 years ago, included information regarding both biotic and abiotic variables, and conducted either statistical or non-statistical analyses related to the functioning of the ecosystem in question.

Information regarding the results obtained during research that satisfied selection criteria was collected and used for identifying patterns, correlations, and mechanisms in the studied phenomena. Various statistical methods reported in the studies (regression analysis, correlation analysis, multivariate statistics) were examined in terms of the interaction between biotic and abiotic variables.

The method makes it possible to conduct a comprehensive comparison between biotic and abiotic interactions in freshwater environments, which helps identify patterns, knowledge gaps, and future research avenues for ecosystem management and conservation.

III. Results

Abiotic Patterns

Abiotic factors, including temperature, hydrology, nutrients, and dissolved oxygen in freshwater habitats, have shown great variability in terms of space and time. There was a flow gradient in rivers and streams that affected the movement of sediments and nutrients, whereas there was temperature and dissolved oxygen stratification in lakes and wetlands, impacting biochemical processes. The concentration of nutrients, especially nitrogen and phosphorus, varied depending on land usage patterns.

Table 1: Abiotic Variables, Associated Biotic Responses, and Ecosystem Function Metrics in Selected Freshwater Systems

Freshwater System	Abiotic Variables (with values)	Biotic Responses	Ecosystem Function Metrics
River	Flow: 0.5–2.0 m/s; Temp: 12–25 °C; DO: 6–10 mg/L; Nitrate: 1–5 mg/L; pH: 6.8–8.2	Fish density: 15–30 ind/m²; Macroinvertebrate richness: 25–45 taxa	Primary productivity: 300–450 mg C/m²/day; Nutrient retention: 60–80%; Decomposition: 1.2–2.0 g/m²/day
Lake	Temp: 10–22 °C (surface); DO: 5–12 mg/L; TP: 20–50 µg/L; Light penetration: 1–5 m	Phytoplankton biomass: 50–200 µg/L; Zooplankton density: 500–1200 ind/L	Nutrient cycling: N 80–95%, P 70–90%; Carbon sequestration: 50–100 g C/m²/year; Productivity: 400–600 mg C/m²/day
Wetland	Water depth: 0.3–1.2 m; Temp: 14–28 °C; pH: 6.5–7.5; Nutrients: N 2–8 mg/L, P 0.1–0.5 mg/L	Macrophyte coverage: 60–90%; Amphibians: 20–35 species; Microbial activity: high	Decomposition: 1.5–3 g/m²/day; N cycling: 70–85%; Organic matter processing: 2–3 kg/m²/year
Floodplain	Flood duration: 5–20 days/year; Sediment: 0.5–2.0 cm deposition; Temp: 15–30 °C	Fish spawning density: 50–200 ind/100 m²; Aquatic plant growth: 60–80% cover	Secondary productivity: 300–500 mg C/m²/day; Nutrient transport: 50–70%
Urban Stream	Flow: 0.3–1.0 m/s; Temp: 15–28 °C; DO: 4–8 mg/L; Pollutants: BOD 3–10 mg/L	Macroinvertebrate richness: 10–20 taxa; Tolerant species dominance	Impaired nutrient cycling: 40–60%; Reduced productivity: 100–250 mg C/m²/day; Decomposition: 0.8–1.5 g/m²/day

Table 1 summarizes important abiotic factors (environmental drivers), for example, temperature, flow rate, dissolved oxygen, and nutrients; it also covers the response to these abiotic drivers by biota in terms

of richness, species composition, and interaction at the trophic level. Some indicators of ecosystem function include the rate of productivity, nutrient turnover, and decomposition.




Figure 2: Relationships between Abiotic Gradients and Biotic Responses Across Freshwater Systems

Figure 2 shows how the important biotic variables, algal biomass, number of plants, number of macroinvertebrates, and fish biomass, are influenced by four principal abiotic factors, including nutrients, temperature, oxygen, and turbidity. Figure 2 (A-D) depicts the effects of variation in environmental parameters on the community structure and productivity, indicating the link between abiotic factors and biotic response.

Biotic Distributions

There was a strong relationship between biodiversity and community structure, with the biotic factors in play. The producer component, comprised of macrophytes and phytoplanktons, was influenced by light, nutrient, and temperature regimes. The consumer groups made up of invertebrates and fish were structured

based on habitat complexities and prey abundance. It was observed through functional group analysis that increased species richness in ecosystems leads to complex food webs.

Interaction Effects

Interactions between the biotic and abiotic elements were made through a number of ways, which helped to define the ecosystem's structure. For example, the provision of nutrients increased the amount of primary production but altered the species composition. The interactions in the food chain were altered. The change in hydrology influenced predation interaction because it altered connectivity between the habitats. Several studies have found that there was a strong correlation between abiotic stressors and community structure alteration.

Ecosystem Functioning Metrics

Functioning of the ecosystem was evaluated through the analysis of various factors such as productivity, nutrient cycling, and decomposition. Species diversity and redundancy ensured nutrient retention and stability of productivity. Wetlands and riparian ecosystems with biotic components had higher levels of decomposition and nitrogen cycling. It indicated that the functioning of ecosystems is governed by the interaction of abiotic and biotic factors.

IV. Discussion

The physical environment, composed of factors like temperature, hydrology, nutrient availability, and oxygen levels, forms one of the basic foundations on which biotic communities can be set up in the freshwater environment. Variation in these physical parameters affects species richness and community composition. For instance, variations in nutrient levels increase the rate of primary production, hence favoring algal and macrophyte proliferation, while variations in the hydrological regime hamper habitat connectivity and the movement of fish. The results corroborate past literature that highlights the significance of environmental heterogeneity in structuring the community in rivers, lakes, and wetlands (Holomuzki et al., 2010).

Biotic interactions such as predation, competition, and mutualism control the ecological functions of the ecosystem (Wohlgemuth et al., 2017). Food chains are complex and redundant, hence increasing stability and ecosystem resilience through effective nutrient recycling and decomposition. Biodiversity is high among the biotic components, and hence, buffering against the adverse effects of the environment (Zivec et al., 2023).

Comparative analysis shows that ecosystems characterized by a high degree of interaction between abiotic and biotic aspects have increased efficiency and resilience (Montoya & Raffaelli, 2010). The review shows that in order to be able to function efficiently, an ecosystem needs bottom-up controls (environmental conditions) and top-down interactions (food chains, trophic levels) (Wang et al., 2022). Literature on wetlands in the tropics, temperate zone river ecosystems, and urban streams demonstrates that environmental conditions and biotic responses play an integral part in influencing ecosystem services (Silknetter et al., 2020; Gerbersdorf et al., 2011).

There are numerous implications for the conservation and management of ecosystems (Barral-Fraga et al., 2020). It is important to conserve natural flow regimes, protect the quality of water, and preserve biodiversity in terms of habitat types in order to sustain a rich community and the operation of the ecosystem itself. Any management practice needs to be concerned with both environmental and biotic factors in order to ensure greater resilience in the face of human-induced stressors and climatic changes.

V. Conclusion

The current review sheds light on the importance of interactions between biotic elements and abiotic factors in determining the structure and functions of freshwater ecosystems. Abiotic factors such as

temperature, hydrology, dissolved oxygen concentration, and nutrients play an important role in controlling species composition, diversity of biotic communities, and trophic interactions within freshwater ecosystems. Conversely, biotic interactions determine ecosystem-level functions including productivity, biogeochemical cycling, decomposition, and energy flow. An analysis of rivers, lakes, wetlands, floodplains, and urban streams suggests that ecosystems with greater functional diversity and complexity in their food webs have better resistance against abiotic stressors. This demonstrates the significance of bottom-up processes, controlled by abiotic factors, and top-down processes, controlled by trophic interactions in driving ecological processes within freshwater ecosystems. The ecological implications of the interaction between abiotic factors and biotic elements include biodiversity conservation, nutrient retention, and ecosystem services. Interruptions in abiotic systems or biotic populations through means of pollution, habitat alteration, and climate change can result in the occurrence of various cascading events that may affect productivity, water quality, and ecosystem resilience. In practice, the management of ecosystem processes requires that natural hydrological processes be maintained, good water quality be preserved, habitat heterogeneity be preserved, and functional biotic communities be restored. Efforts towards ecosystem conservation and restoration require that both abiotic and biotic factors be considered as an integrated system that builds up resilience and buffers against disturbances, thus enabling the continuous provision of critical ecosystem services.

References

[1] Holomuzki, J. R., Feminella, J. W., & Power, M. E. (2010). Biotic interactions in freshwater benthic habitats. *Journal of the North American Benthological Society*, *29*(1), 220-244. https://doi.org/10.1899/08-044.1

[2] Vaughn, C. C. (2010). Biodiversity losses and ecosystem function in freshwaters: emerging conclusions and research directions. *BioScience*, *60*(1), 25-35. https://doi.org/10.1525/bio.2010.60.1.7

[3] Manthila, P., Anuradha, K. M., & Chandrakumar, R. (2025). Climate-Driven Alterations in Aquatic Biodiversity and Ecosystem Functioning Across Freshwater and Coastal Systems. *Journal of Aquatic Ecology and Environmental Sustainability*, 2(3), 26-34.

[4] Atkinson, C. L., Capps, K. A., Rugenski, A. T., & Vanni, M. J. (2017). Consumer-driven nutrient dynamics in freshwater ecosystems: From individuals to ecosystems. *Biological Reviews*, *92*(4), 2003-2023.https://doi.org/10.1111/brv.12318

[5] Silknetter, S., Creed, R. P., Brown, B. L., Frimpong, E. A., Skelton, J., & Peoples, B. K. (2020). Positive biotic interactions in freshwaters: A review and research directive. *Freshwater Biology*, *65*(4), 811-832. https://doi.org/10.1111/fwb.13476

[6] Zivec, P., Sheldon, F., & Capon, S. J. (2023). Natural regeneration of wetlands under climate change. *Frontiers in Environmental Science*, *11*, 989214. https://doi.org/10.3389/fenvs.2023.989214

[7] Setubal, R. B., Petry, A. C., Bonecker, C. C., Martins, T., Nova, C. C., Figueiredo-Barros, M. P., & Bozelli, R. L. (2020). Biotic factors determine ecosystem processes in environments with different hydrological regimes. *Freshwater Biology*, *65*(8), 1376-1391. https://doi.org/10.1111/fwb.13506

[8] Gerbersdorf, S. U., Hollert, H., Brinkmann, M., Wieprecht, S., Schüttrumpf, H., & Manz, W. (2011). Anthropogenic pollutants affect ecosystem services of freshwater sediments: the need for a “triad plus x” approach. *Journal of Soils and Sediments*, *11*(6), 1099-1114. https://doi.org/10.1007/s11368-011-0373-0

[9] Montoya, J. M., & Raffaelli, D. (2010). Climate change, biotic interactions and ecosystem services. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1549), 2013. https://doi.org/10.1098/rstb.2010.0114

[10] Barral-Fraga, L., Barral, M. T., MacNeill, K. L., Martiñá-Prieto, D., Morin, S., Rodríguez-Castro, M. C., ... & Guasch, H. (2020). Biotic and abiotic factors influencing arsenic biogeochemistry and

toxicity in fluvial ecosystems: a review. *International journal of environmental research and public health*, *17*(7), 2331. https://doi.org/10.3390/ijerph17072331

[11] Nõges, P., Argillier, C., Borja, Á., Garmendia, J. M., Hanganu, J., Kodeš, V., ... & Birk, S. (2016). Quantified biotic and abiotic responses to multiple stress in freshwater, marine and ground waters. *Science of the Total Environment*, *540*, 43-52. https://doi.org/10.1016/j.scitotenv.2015.06.045

[12] Wohlgemuth, D., Solan, M., & Godbold, J. A. (2017). Species contributions to ecosystem process and function can be population dependent and modified by biotic and abiotic setting. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1855), 20162805. https://doi.org/10.1098/rspb.2016.2805

[13] Birk, S., Chapman, D., Carvalho, L., Spears, B. M., Andersen, H. E., Argillier, C., ... & Hering, D. (2020). Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems. *Nature Ecology & Evolution*, *4*(8), 1060-1068. https://doi.org/10.1038/s41559-020-1216-4

[14] Rodrigues-Filho, J. L., Macêdo, R. L., Sarmento, H., Pimenta, V. R., Alonso, C., Teixeira, C. R., ... & Cionek, V. M. (2023). From ecological functions to ecosystem services: linking coastal lagoons biodiversity with human well-being. *Hydrobiologia*, *850*(12), 2611-2653. https://doi.org/10.1007/s10750-023-05171-0

[15] Wang, J., Hu, A., Meng, F., Zhao, W., Yang, Y., Soininen, J., ... & Zhou, J. (2022). Embracing mountain microbiome and ecosystem functions under global change. *New Phytologist*, *234*(6), 1987-2002. https://doi.org/10.1111/nph.18051