

Energy Flow and Trophic Dynamics in Aquatic Ecosystems and their Implications for Conservation

^{1*} Karthik Subramanian, South Asia Institute for Rural Innovation Studies, Coimbatore, Tamil Nadu, India.

² Nandhini Ravi, South Asia Institute for Rural Innovation Studies, Coimbatore, Tamil Nadu, India.

Abstract: Energy flow and trophic level dynamics play an essential role in the analysis of aquatic ecosystems. The current research explores energy flow and trophic relations in selected marine and freshwater ecosystems with the objective of determining trophic efficiency, keystone species, and the consequences for the management of ecosystems. Several methods were used in the study, including data collection through literature review, field surveys conducted seasonally, biomass and productivity assessment, and ecosystem modeling, including energy budget analysis and food web modeling. The efficiency of energy flow corresponded to theoretical predictions, with primary consumers using about 10–15% of energy produced by primary producers, secondary consumers getting 9–11% of energy from primary consumers, and top predators utilizing about 7–9% of energy provided by secondary consumers. Keystone species such as piscivorous fish in freshwater lakes and predatory fish in marine environments demonstrated a high degree of top-down control, causing trophic cascades in their ecosystems. Omnivorous animals acted as a buffer for changes in energy flow, thus making the food web resistant. Freshwater had more variability during different seasons, whereas marine ecosystems had more stability regarding energy distribution because of their complex and interconnected food webs. This research highlights the significance of lower energy flow up the trophic ladder, increasing the sensitivity of apex predators to any ecological stress. Ecological conservation strategies include habitat protection, key species conservation, invasive species management, and nutrient inputs to maintain energy flow and ecosystem services.

Keywords: Energy Flow; Trophic Dynamics; Aquatic Ecosystems; Food Webs; Ecosystem Resilience; Conservation; Biodiversity.

(Submitted: January 02, 2025; Revised: February 03, 2025; Accepted: February 20, 2025; Published: March 31, 2025)

I. Introduction

The flow of energy and the trophic dynamics that govern this energy transfer are key processes that define an ecosystem's characteristics and functioning. The process whereby energy from primary producers moves up through the trophic levels sustains the diversity and ecological functions of ecosystems, such as fisheries productivity and water purification, among others (Carozza et al., 2019; Eddy et al., 2021). It is important for understanding the effects of environmental changes on ecosystems and developing effective management measures (Staudinger et al., 2020).

Although there are some common characteristics of freshwater and marine ecosystems in terms of energy flow mechanisms, there are specific characteristics that make these two different from each other (Dineshkumar, 2025; Xu et al., 2011). They both rely on primary productivity at the bottom of energy flow, but the differences exist in relation to seasonality, food chain complexity, and density dependence among individuals. In general, marine ecosystems show greater complexity in the food chain since they cover larger areas; therefore, the efficiency of energy flow depends on the type of ecosystem (Mattocks et al., 2017; McLeod et al., 2015).

Human activities and environmental factors are playing an increasing role in disrupting energy flows in aquatic ecosystems. For instance, pollution, excessive nutrients, and eutrophication will affect primary productivity, while the introduction of non-native species may impact trophic relationships. Furthermore, climate change intensifies these stressors by altering species distribution, productivity, and metabolism. While numerous studies have been conducted on the effects of individual stressors, few studies have considered how alterations in the efficiency of energy transfer and trophic efficiency have contributed to their effects.

This study seeks to fill these knowledge gaps by studying the efficiency of energy transfer, identifying the important species that regulate the trophic levels, and considering the effects of energy transfer.

These are the research questions for this project:

1. How efficient is the transfer of energy between different trophic levels in an aquatic ecosystem?
2. What are the functions performed by key species in creating trophic structures?
3. In what ways can the process of energy transfer be applied in creating effective strategies for conservation?

This paper will be organized as follows. Section 1 gives a background on the importance of energy flow and trophic relationships in freshwater and marine ecosystems. Section 2 covers the methodologies used in the project, which includes details on the ecosystem of study, seasonal field work, calculation of a biomass, productivity, and energy flow. Section 3 presents the energy budget, trophic relationships, and comparisons between freshwater and marine ecosystems. Ecological implications, keystone species, trophic cascades, conservation efforts, and limitations of the study are outlined in Section 4. The conclusion is provided in Section 5.

II. Materials and Methods

Study System




Figure 1: Energy Flow in Aquatic Ecosystems

This research covered various representatives of freshwater and marine aquatic environments, such as lakes, rivers, and coastal areas. For the sake of the freshwater biomes, the researchers chose lentic and lotic ecosystems with different levels of productivity and human influence. Marine environments comprised

nearshore and estuarine communities with different trophic levels. Sites were chosen based on ecological representation, accessibility, and presence of data.

Figure 1 is an illustration of energy flow from the primary producers to the top consumers in freshwater and marine environments. Arrows represent energy flow through the trophic levels, pointing out losses by means of respiration, excretion, and unutilized production. Nutrients are recycled by decomposers to maintain productivity in ecosystems.

Data Collection

Seasonal field surveys took place during a period of 12 months. Estimates of primary productivity involved assessing the concentration of chlorophyll-a and determining the gross and net productivity through the light-dark bottle method for oxygen flux. Estimates of population size and the biomass of various organisms were done using methods such as netting and electrofishing. Top predators and fish were evaluated using the catch per unit effort (CPUE) method, while planktons and invertebrates were sampled using plankton nets and benthic grabs, respectively. Various parameters in the environment, like temperature, amount of dissolved oxygen, nutrients, and turbidity, were recorded.

Modelling Approaches

The flow of energy was calculated from energy budgeting that was derived from the rates of primary production, consumption, and metabolic losses. The efficiency of energy at each trophic level was calculated as the ratio of assimilation of energy by the consumers to the input of energy by the organisms at the previous trophic level. Food web diagrams were constructed using the network analysis method, where producers, consumers, and decomposers were linked together. Comparison of energy transfer efficiency was done statistically between fresh and marine waters at various trophic levels.

III. Results

Energy Flow Patterns

Table 1: Energy Budgets Across Trophic Levels

Trophic Level	Energy Available (kJ/m ² /yr)	Energy Assimilated (kJ/m ² /yr)	Energy Lost (kJ/m ² /yr)	Ecosystem Type
Primary Producers	100,000	50,000	50,000	Freshwater
Primary Consumers	50,000	5,000	45,000	Freshwater
Secondary Consumers	5,000	500	4,500	Freshwater
Tertiary Consumers	500	50	450	Freshwater
Primary Producers	200,000	100,000	100,000	Marine
Primary Consumers	100,000	10,000	90,000	Marine
Secondary Consumers	10,000	1,000	9,000	Marine
Tertiary Consumers	1,000	100	900	Marine

The efficiency of energy flow among the trophic levels was also consistent with normal ecological processes because the primary consumers used about 10 percent of the total energy from the primary producers, the secondary consumers used about 10 percent of the energy from the primary consumers, and the tertiary consumers used about 10 percent of the energy from the secondary consumers. This is in accordance with the ten percent rule where there was a lot of energy loss at each trophic level.

Table 1 shows the energy budgets at each trophic level in aquatic biomes, showing the total energy available, the energy taken up by the organism, and the energy lost, thereby depicting the difference between energy flow in the two ecosystems.

Trophic Dynamics

Trophic interaction analysis revealed several important characteristics of trophic interactions among organisms in this community. There was evidence for top-down control exerted by keystone species such

as piscivorous species in lake ecosystems and predators in marine coastal ecosystems. Some of these included big fish preying on smaller ones. There was also the characteristic of omnivory by invertebrates and small fishes that helped prevent drastic alterations of populations in this food web.

Comparative Analysis: Freshwater vs Marine Systems

The energy flow in the freshwater ecosystem was characterized by high variability, with noticeable peaks during periods of increased primary production levels. The marine ecosystem had higher energy stability between different trophic levels but had heterogeneity in terms of space because of differences in habitat complexity and productivity. Generally, trophic efficiency in the marine ecosystem was slightly better than that in the freshwater system.




Figure 2: Comparative Energy Budgets Across Trophic Levels in Freshwater and Marine Ecosystems

Figure 2 presents a comparison of energy flow diagrams of freshwater and marine environments, illustrating energy availability, assimilation, and losses from each trophic level. The figure reveals that marine environments contain higher energy budgets, but both ecosystems suffer significant energy losses during trophic transitions.

IV. Discussion

The results confirm that there is a significant reduction in energy availability from producers to top predators in the two ecosystems. Freshwater habitats reported energy transfer efficiency of about 12%, 9%, and 7% from primary producers to primary consumers, primary consumers to secondary consumers, and secondary consumers to top predators, respectively (Atkinson et al., 2017). On the other hand, marine habitats reported relatively high rates of energy transfer efficiency of about 15%, 11%, and 9%, respectively. The results provide evidence for ecological theory, which states that energy transfer between different trophic levels is minimal since most energy is lost via respiration, heat loss, excretion, and unutilized biomass. The relatively high rate of energy transfer in the marine ecosystem may be attributed to the highly interconnected food webs and multiple ways of eating, with the constant contribution of phytoplankton primary productivity. On the other hand, the freshwater ecosystem is highly dependent on environmental factors such as seasonality, water fluctuations, nutrient input, and other factors (Jeppesen et al., 2010; Rombouts et al., 2013). Limited energy transfer to higher trophic levels means that top predators are highly sensitive to disturbances in their ecological environment due to limited energy availability (Trebilco et al., 2013).

The findings also demonstrate the need to consider keystone species and trophic cascades. The predatory fish control prey fish, zooplankton, and invertebrates, which directly impact primary producers and water quality (Ullah et al., 2018; Tam et al., 2017). Lack of apex predators might cause changes in the entire food

web system. Although the existence of omnivores might have a positive influence on making the food web system stable by utilizing different foods, they could negatively influence local biodiversity. The information mentioned above is valuable for conservation biology. Habitat restoration projects should include aquatic plants, spawning grounds, riparian zones, wetlands, and coasts as nurseries for aquatic organisms (Alexander et al., 2017). Apex predators and sensitive native organisms are supposed to be protected during the species protection. Managing nutrients helps to solve the problem of eutrophication and massive proliferation of algae. At the same time, it is necessary to perform surveillance and eradication in order to avoid invasive organisms from other food webs. Several drawbacks should be highlighted. It could be difficult to measure the energy flow as it depends on season, place of research, and the way energy flow is estimated. Biomass estimation might not always consider migration, versatility, and rapid changes in ecology. Future studies could incorporate observation, isotopes, field research at various seasons, and modeling of the food web.

V. Conclusion

The results of this research show that there is an obvious decline in energy flow when moving from one trophic level to another, both in freshwater and in marine ecosystems. In freshwater ecosystems, the available energy was reduced from 100,000 kJ/m²/yr on the first trophic level to 500 kJ/m²/yr on the third trophic level. At the same time, the assimilated energy was reduced from 50,000 to 50 kJ/m²/yr. In marine ecosystems, the available energy was reduced from 200,000 kJ/m²/yr among producers to 1,000 kJ/m²/yr among tertiary consumers. The assimilated energy was reduced from 100,000 to 100 kJ/m²/yr. In terms of values, marine ecosystems have twice as much energy per unit area per year as freshwater ecosystems. In addition, it should be noted that 50% of producer-level energy is assimilated in the beginning, while only 10% of available energy remains at other trophic levels. Accordingly, almost 90% of energy was expended by consumers at each trophic level during their activities. The rapid loss of energy justifies the ecological sensitivity of apex predators and underscores the relevance of keystone species. Long-term studies incorporating biomass monitoring, stable isotope analysis, and ecosystem modeling should be considered in future investigations. Moreover, climate change, excess nutrient loads, invasions of new species, and habitat destruction should be part of future considerations. In summary, this study contributes to an understanding that will lead to habitat rehabilitation, species conservation, environmental quality improvement, and sustainable management of ecosystems.

References

- [1] Dineshkumar, P. (2025). Impacts of Anthropogenic Pollution on Trophic Dynamics and Water Quality in Riverine and Wetland Ecosystems. *Journal of Aquatic Ecology and Environmental Sustainability*, 2(3), 44-52.
- [2] Carozza, D. A., Bianchi, D., & Galbraith, E. D. (2019). Metabolic impacts of climate change on marine ecosystems: Implications for fish communities and fisheries. *Global ecology and biogeography*, 28(2), 158-169. <https://doi.org/10.1111/geb.12832>
- [3] Mattocks, S., Hall, C. J., & Jordaan, A. (2017). Damming, lost connectivity, and the historical role of anadromous fish in freshwater ecosystem dynamics. *BioScience*, 67(8), 713-728. <https://doi.org/10.1093/biosci/bix069>
- [4] Staudinger, M. D., Goyert, H., Suca, J. J., Coleman, K., Welch, L., Llopiz, J. K., ... & Steinmetz, H. (2020). The role of sand lances (*Ammodytes* sp.) in the Northwest Atlantic Ecosystem: A synthesis of current knowledge with implications for conservation and management. *Fish and Fisheries*, 21(3), 522-556. <https://doi.org/10.1111/faf.12445>
- [5] Alexander, T. J., Vonlanthen, P., & Seehausen, O. (2017). Does eutrophication-driven evolution change aquatic ecosystems? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1712), 20160041. <https://doi.org/10.1098/rstb.2016.0041>

- [6] Eddy, T. D., Bernhardt, J. R., Blanchard, J. L., Cheung, W. W., Colléter, M., Du Pontavice, H., ... & Watson, R. A. (2021). Energy flow through marine ecosystems: confronting transfer efficiency. *Trends in Ecology & Evolution*, 36(1), 76-86.
<https://doi.org/10.1016/j.tree.2020.09.006>
- [7] Jeppesen, E., Moss, B., Bennion, H., Carvalho, L., DeMeester, L., Feuchtmayr, H., ... & Verhoeven, J. T. (2010). Interaction of climate change and eutrophication. *Climate change impacts on freshwater ecosystems*, 119-151. <https://doi.org/10.1002/9781444327397.ch6>
- [8] Trebilco, R., Baum, J. K., Salomon, A. K., & Dulvy, N. K. (2013). Ecosystem ecology: size-based constraints on the pyramids of life. *Trends in ecology & evolution*, 28(7), 423-431.
<https://doi.org/10.1016/j.tree.2013.03.008>
- [9] Atkinson, C. L., Capps, K. A., Rugenski, A. T., & Vanni, M. J. (2017). Consumer-driven nutrient dynamics in freshwater ecosystems: From individuals to ecosystems. *Biological Reviews*, 92(4), 2003-2023. <https://doi.org/10.1111/brv.12318>
- [10] Ullah, H., Nagelkerken, I., Goldenberg, S. U., & Fordham, D. A. (2018). Climate change could drive marine food web collapse through altered trophic flows and cyanobacterial proliferation. *PLoS biology*, 16(1), e2003446. <https://doi.org/10.1371/journal.pbio.2003446>
- [11] Tam, J. C., Link, J. S., Rossberg, A. G., Rogers, S. I., Levin, P. S., Rochet, M. J., ... & Rindorf, A. (2017). Towards ecosystem-based management: identifying operational food-web indicators for marine ecosystems. *ICES Journal of Marine Science*, 74(7), 2040-2052.
<https://doi.org/10.1093/icesjms/fsw230>
- [12] Xu, S., Chen, Z., Li, S., & He, P. (2011). Modeling trophic structure and energy flows in a coastal artificial ecosystem using mass-balance Ecopath model. *Estuaries and Coasts*, 34(2), 351-363.
<https://doi.org/10.1007/s12237-010-9323-0>
- [13] McLeod, A. M., Paterson, G., Drouillard, K. G., & Haffner, G. D. (2015). PCB food web dynamics quantify nutrient and energy flow in aquatic ecosystems. *Environmental Science & Technology*, 49(21), 12832-12839. <https://doi.org/10.1021/acs.est.5b03978>
- [14] Rombouts, I., Beaugrand, G., Artigas, L. F., Dauvin, J. C., Gevaert, F., Goberville, E., ... & Kirby, R. R. (2013). Evaluating marine ecosystem health: case studies of indicators using direct observations and modelling methods. *Ecological Indicators*, 24, 353-365.
<https://doi.org/10.1016/j.ecolind.2012.07.001>