

Human Impact on the Hydrological Cycle and Consequences for Aquatic Ecosystem Services

^{1*} Karthik Subramanian, South Asia Institute for Rural Innovation Studies, Coimbatore, Tamil Nadu, India.

² Nandhini Ravi, South Asia Institute for Rural Innovation Studies, Coimbatore, Tamil Nadu, India.

Abstract: The hydrological cycle is essential to a stable global climate and healthy ecosystems, yet it is rapidly changing due to human actions. Human activities, including urban sprawl, industrialized agriculture, large-scale dam construction, and groundwater overdraft, have dramatically altered the natural flow regime, sediment delivery, and water quality. Although numerous studies have explored human-induced hydrological changes individually, there remains a disconnect between quantitatively modeling hydrological processes and evaluating fundamental ecosystem services (water purification, flood pulse management, habitat integrity). In this paper, it analyzes existing data on human impacts on the hydrological cycle and describe the cascade of socio-ecological impacts on the coastal wetland-riverine ecotone of the Ganges-Brahmaputra Delta. It identifies how flow regulation and increased intensity of land use degrade ecological connectivity and lead to fragmented habitats, shifted salinity regimes, eutrophication, and degraded trophic structure, resulting in a decrease in provision, regulation, and cultural services. Maladaptive feedback is evident: as natural processes decline, it relies on brittle 'grey' infrastructure, which further degrades hydrology and degradation; only by merging nature-based restoration with new technological tools like remote sensing and decentralized AI-driven predictive models will it establish restorative resilience and ensure water security.

Keywords: Hydrological Cycle; Ecosystem Services; Anthropogenic Impact; Ganges-Brahmaputra Delta; Nature-Based Solutions; Remote Sensing; Ecological Resilience.

(Submitted: March 28, 2025; Revised: April 15, 2025; Accepted: May 27, 2025; Published: June 30, 2025)

I. Introduction

The hydrologic cycle, a continuously active and fundamentally significant process driven by solar energy and gravity, circulates water among the atmosphere, land surface, and subsurface, governing climate, regulating life, and ultimately mediating all of the planet's surface processes through precipitation, evaporation, transpiration, infiltration, and runoff. The naturally functioning circulatory process is a master distributor of nutrients and energy throughout the planet and is the driving force behind healthy and stable aquatic ecosystems worldwide.

The unprecedented speed of human endeavors in the last hundred years has irrevocably perturbed this system, transforming natural rates and routes of water movement through increased urbanization, intensive agriculture, dam construction, and heavy groundwater withdrawal (Smale et al., 2019). The physical disruption of natural drainage systems, heightened pollution, and changes to seasonal flow patterns have altered the historically linked system and driven aquatic ecosystems into state changes through which their function is greatly diminished (Patel, 2024; Terrado et al., 2014; Buytaert et al., 2011).

Though substantial data exists concerning separate water resource processes, these studies are disparate and provide no synthesized picture of how disparate changes in these processes translate into impaired ecosystem services (Yang et al., 2021; Gordon et al., 2010; El Jeitany et al., 2024). The gap between quantitative modeling of hydrologic processes and the social/economic assessment of individual services (e.g., Natural water purification, maintenance of flood pulses, and support of fish/wildlife habitat) is currently immense, but must be addressed for us to move from descriptive science toward predictive and proactive management for the water resources.

Objective

- To synthesize the current evidence of how exactly humans are changing hydrology.
- To assess the cascading socio-ecological implications of these changes in the provision of key aquatic ecosystem services.
- To suggest a complete scheme for coastal wetland-riverine interface systems in the Ganges-Brahmaputra Delta that connects hydrological sustainability with essential ecosystem functions.

Logical progression of topics, including a section each for hydrological modifications and ecological response. Hydrological modifications, detailing changes brought by urbanization, dam building, agriculture, and groundwater depletion, and accompanied by a conceptual diagram (section II), and Ecological Response detailing changes to habitats, salinity, and the trophic web (section III) are separated out for clarity. The results are broken down into a discussion of ecosystem services, namely provisioning, regulating, and cultural (section IV, presented with a summary table), and synthesis and discussion focusing on maladaptive feedback loops and the integration of artificial intelligence with nature-based approaches (section V). Section VI presents the conclusion and relevant policy recommendations.

II. Mechanisms of Human Impact

Hydrological Alteration

Human activities have profoundly changed the hydrology of the Ganges-Brahmaputra deltaic systems by modifying flow patterns and intensive land use. Dams and river diversion upstream have reduced the natural seasonal floods that once coordinated sediment delivery and water supply throughout the system. Intensifying urbanization and development have made flash flood peaks larger and more frequent, and have reduced response times to peak storm discharge. This flashy flow regime is compounded by persistent agricultural input of nitrogen and phosphorus, changing the nutrient mass balance and lowering the ability of the system to flush during the dry season. Groundwater extraction, mostly for agriculture, is contributing to land subsidence and base-flow depletion; consequently, the freshwater-seawater interface is moving landward due to salinity intrusion.

Ecological Response

These changes in hydrological regime have serious impacts on the ecological integrity of aquatic ecosystems by subjecting them to new physical conditions, thus forcing the ecological systems to evolve to the new stimuli. Attenuation of seasonal flood pulse resulted in a reduction of connectivity between the main river channel and flood plains and consequent loss of spawning and nursery habitats of migratory fish, which need these habitats for successful recruitment. Altered salinity gradient caused by reduced freshwater inflow and land subsidence caused destruction of salt-sensitive vegetation and invasion of brackish species, hence altering the composition of the benthic community structure. Nutrient enrichment coupled with a reduction in flow velocity causes proliferation of toxic algal blooms, which reduces the dissolved oxygen concentration, causes hypoxic areas, and subsequently undermines all aspects of the trophic web by damaging producers and important invertebrates.

III. Consequences for Ecosystem Services

Provisioning Services

The modification of the hydrological cycle jeopardizes the local resource base on which livelihoods depend primarily by diminishing the inland and estuarine fishery resources. Change of seasonal flow regimes, coupled with the invasion of saline waters, limits the freshwater supply to irrigation, thereby decreasing crop yields in the deltaic region. Reduced availability of terrestrial resources like timber,

herbaceous plant materials from river flood plains, and within the river limits, access for communities to collect.

Table 1: Quantitative Summary of Human Activities and Associated Ecosystem Service Impacts

Human Activity	Hydrological Change	Ecological Consequence	Ecosystem Services Affected	Severity
Urbanization	Increased Impervious Surfaces, Altered Runoff	Habitat Fragmentation, Reduced Connectivity	Provisioning, Regulating, Cultural	High
Dam Construction	Flow Regulation, Seasonal Flow Alteration	Altered Flood Pulses, Ecosystem Disruption	Provisioning, Regulating	High
Agriculture	Nutrient Loading, Irrigation Runoff	Algal Blooms, Hypoxia, Degraded Water Quality	Provisioning, Regulating	Medium
Groundwater Extraction	Declining Groundwater Table, Reduced Baseflow	Saltwater Intrusion, Lower Dry-Season Flows	Provisioning, Regulating	High
Land Use Change	Deforestation, Wetland Conversion	Loss of Biodiversity, Altered Community Composition	Provisioning, Cultural	Medium

Table 1 summarizes in simple terms the main human activities that affect the hydrological cycle and the resulting environmental impacts, showing a correlation between an activity, hydrologic changes resulting from it, and the measured ecological impacts. Ecosystem services impacted, such as provisioning (fishing and irrigation uses), regulating (flood control, water purification uses), and cultural ones, and intensity ratings show relative impact level from each of the activities, enabling visual comparison.




Figure 1: Seasonal Variation in Ecosystem Service Indicators under Anthropogenic Pressure

Shows figure 1 months/seasons, Y-axis presents normalized indicators such as freshwater availability, dissolved oxygen, and habitat connectivity; each ecosystem service (provisioning, regulating, cultural) is presented with a number of lines/bars. Shows how services decrease during high impacts of human activity (peak irrigation, urban runoff, etc.).

Regulating Services

Anthropogenic changes to hydrology strongly damage the ecosystem's ability to deliver key regulatory services. With the natural floodplain connection lost, the landscape is less able to dampen extreme floods, and downstream communities become even more susceptible to extreme weather. Meanwhile, the reduced natural filtration of water and impaired natural nutrient cycling lessen the ability of the ecosystem to process and remove from the environment pollutants found in water, thus decreasing water quality overall. Furthermore, the disappearance of coastal wetland and mangrove habitats-resulting from the disruption of

natural salinity gradients and subsidence-removes the ecosystem's natural capacity to sequester carbon, instead transforming carbon sinks into sources of greenhouse gas.

Cultural/Supporting Services

The ecological degradation extends to impacts on cultural and supporting services, thereby changing the relationship between local communities and their aquatic ecosystems. Reduced biodiversity, particularly iconic species and keystone species, reduces the natural value of the ecosystem and consequently decreases the opportunities for eco-tourism and recreation that rely on pristine environments. It further damages cultural practice and traditional knowledge that is intrinsically tied to the historic hydrological regimes of the Ganges-Brahmaputra Delta, representing a loss of socio-ecological heritage.

IV. Synthesis and Discussion

Ecological degradation of aquatic ecosystem services often initiates a maladaptive feedback cycle whereby the ecosystem's loss of natural capacity results in intensified anthropogenic intervention (Pataki et al., 2011). As natural regulation capacity is degraded by changing hydrological regimes, there is socio-economic pressure to engineer 'grey infrastructure' (embankments, levees) to mitigate direct impacts to downstream urban and agricultural regions. Rigid grey infrastructure only intensifies hydrological instability by creating stronger disconnection between riverine and floodplain environments. This loss of connectivity only accelerates ecological degradation by further impacting natural filtering capacity and causing the loss of crucial aquatic habitat. This ultimately fosters dependence on high-maintenance, non-renewable infrastructure instead of naturally occurring, renewable, self-sustaining ecosystem services (Sohoulande Djebou & Singh, 2016; Livesley et al., 2016).

The cycle must be broken by pairing innovative technological monitoring with a restoration-based management approach. The use of high-resolution remote sensing data combined with decentralized artificial intelligence inference can monitor hydrological stressors, such as salinity gradients and the rate of land subsidence, on the landscape and at a fine resolution in the deltaic regions. Hybrid machine learning models (such as CNN-LSTMs) can provide prognoses of the impacts that extreme weather will have on aquatic ecosystem service delivery and allow for proactive resource management (Talbot et al., 2018). Reestablishing mangrove and tidal wetland habitats will then allow for ecosystem-based defenses against storm surge and enhanced carbon sequestration (Grizzetti et al., 2016; Vigerstol & Aukema, 2011). By linking these solutions to ecosystem service provision, managers can begin to balance anthropogenic demands (industrial and agricultural) with those that will preserve natural water security and biodiversity (Rocha et al., 2015).

V. Conclusion and Policy Recommendations

This study clearly confirms that changes to hydrological regimes are an existential threat to the productive capacity and integrity of the Ganges-Brahmaputra deltaic system. The loss of natural hydrological dynamics as a result of unabated urbanization, targeted damming of river systems, and unsustainable abstraction rates initiates a cascading ecological decline that impacts a wide range of provisioning, regulating, and cultural ecosystem services. Management practices that exclusively favor the implementation of fixed, "grey" engineering solutions are inherently counterproductive, as they lock us into a vicious cycle of degradation and reliance on increasingly costly, non-renewable technologies that disconnect us further from the dynamic hydrological coupling of river and floodplain. Policies must be urgently redirected toward adaptive, scientific solutions that re-establish natural hydrological connectivity. Stakeholders in the region must embrace Nature-Based Solutions (NbS) like mangrove reforestation and floodplain reconnection as primary defenses against environmental stressors and restore filtering and carbon sequestration capabilities. These must be coupled with modern data-driven monitoring using high-resolution remote sensing and decentralized, artificial intelligence-powered predictive modeling to support truly adaptive resource management. Finally, future research efforts must focus on the

quantification of the ecological service trade-offs inherent in these endeavors, clearly linking water security with broader regional socio-economic health. Only by re-establishing strict adherence to hydrological system limits can it shift from crisis mitigation to proactive, restorative resilience.

References

- [1] Patel, P. (2024). GIS-Based Urban Flood Prediction and Drainage System Optimization Using Machine Learning Techniques. *Transactions on Secure Communication Networks and Protocol Engineering*, 27-32.
- [2] Talbot, C. J., Bennett, E. M., Cassell, K., Hanes, D. M., Minor, E. C., Paerl, H., ... & Xenopoulos, M. A. (2018). The impact of flooding on aquatic ecosystem services. *Biogeochemistry*, 141(3), 439-461. <https://doi.org/10.1007/s10533-018-0449-7>
- [3] Yang, D., Yang, Y., & Xia, J. (2021). Hydrological cycle and water resources in a changing world: A review. *Geography and sustainability*, 2(2), 115-122. <https://doi.org/10.1016/j.geosus.2021.05.003>
- [4] Terrado, M., Acuña, V., Ennaanay, D., Tallis, H., & Sabater, S. (2014). Impact of climate extremes on hydrological ecosystem services in a heavily humanized Mediterranean basin. *Ecological indicators*, 37, 199-209. <https://doi.org/10.1016/j.ecolind.2013.01.016>
- [5] Gordon, L. J., Finlayson, C. M., & Falkenmark, M. (2010). Managing water in agriculture for food production and other ecosystem services. *Agricultural Water Management*, 97(4), 512-519. <https://doi.org/10.1016/j.agwat.2009.03.017>
- [6] Rocha, J., Yletyinen, J., Biggs, R., Blenckner, T., & Peterson, G. (2015). Marine regime shifts: drivers and impacts on ecosystems services. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370(1659), 20130273. <https://doi.org/10.1098/rstb.2013.0273>
- [7] Grizzetti, B., Lanzaova, D., Lique, C., Reynaud, A., & Cardoso, A. C. (2016). Assessing water ecosystem services for water resource management. *Environmental Science & Policy*, 61, 194-203. <https://doi.org/10.1016/j.envsci.2016.04.008>
- [8] El Jeitany, J., Pacetti, T., & Caporali, E. (2024). Evaluating climate change effects on hydrological functionality and water-related ecosystem services. *Ecohydrology*, 17(4), e2557. <https://doi.org/10.1002/eco.2557>
- [9] Buytaert, W., Cuesta-Camacho, F., & Tobón, C. (2011). Potential impacts of climate change on the environmental services of humid tropical alpine regions. *Global Ecology and Biogeography*, 20(1), 19-33. <https://doi.org/10.1111/j.1466-8238.2010.00585.x>
- [10] Pataki, D. E., Carreiro, M. M., Cherrier, J., Grulke, N. E., Jennings, V., Pincetl, S., ... & Zipperer, W. C. (2011). Coupling biogeochemical cycles in urban environments: ecosystem services, green solutions, and misconceptions. *Frontiers in Ecology and the Environment*, 9(1), 27-36. <https://doi.org/10.1890/090220>
- [11] Smale, D. A., Wernberg, T., Oliver, E. C., Thomsen, M., Harvey, B. P., Straub, S. C., & Moore, P. J. (2019). Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nature Climate Change*, 9(4), 306-312. <https://doi.org/10.1038/s41558-019-0412-1>
- [12] Vigerstol, K. L., & Aukema, J. E. (2011). A comparison of tools for modeling freshwater ecosystem services. *Journal of environmental management*, 92(10), 2403-2409. <https://doi.org/10.1016/j.jenvman.2011.06.040>
- [13] Sohoulade Djebou, D. C., & Singh, V. P. (2016). Impact of climate change on the hydrologic cycle and implications for society. *Environment and Social Psychology*, 1(1), 36-49. <https://doi.org/10.18063/esp.2016.01.002>
- [14] Livesley, S. J., McPherson, E. G., & Calfapietra, C. (2016). The urban forest and ecosystem services: impacts on urban water, heat, and pollution cycles at the tree, street, and city scale. *Journal of environmental quality*, 45(1), 119-124. <https://doi.org/10.2134/jeq2015.11.0567>