Aquatic Ecosystems and Environmental Frontiers, Volume 3, Issue 2, June 2025 *ISSN: 3049-186X*
https://doi.org/10.70102/AEEF/V3I2/14

Integrating Hydrology and Ecosystem Functioning in Water Resource Management

[1*] Mehmet Yılmaz, Anatolian Institute for Advanced Technology Studies, Ankara, Turkey.

[2] Elif Kaya, Anatolian Institute for Advanced Technology Studies, Ankara, Turkey.

Abstract: The history of water resource management is marked by an engineering-centered paradigm that, in its focus on anthropogenic usefulness through the mechanistic control of flow regimes, has disregarded the fundamental requirements of biological components in aquatic systems. The resulting heritage of siloed water governance has led to widespread ecological degradation: lost hydrological connections, reduced habitat availability, and interrupted nutrient cycles. This paper integrates the current scientific knowledge of ecohydrological processes with the need to connect physical hydrological models with biological indicators of performance. It draws upon an analysis of feedback loops between flow, groundwater baseflow, and ecosystem processes to describe how the non-linear responses of biological communities to hydrological manipulation jeopardize long-term water system resilience. It offers an Integrated Eco-Governance Framework (IEGF) to address these pervasive problems beyond the static approach of allocation frameworks. IEGF builds on three new principles: (1) the introduction of dynamic functional flow regimes that mimic natural hydrologic variability, (2) the establishment of cooperative benefit-sharing arrangements for the co-management of water resources among multiple stakeholders, and (3) the incorporation of adaptive management through real-time, on-the-ground indicators of ecosystem integrity. IEGF aims to transform human relations with water from commodity use to a system management context that honors the functioning ecological system in which water exists. The inclusion of ecological integrity as a primary technical design parameter enables policymakers to balance human needs for water security with ecological demands for essential ecosystem services and resilience to hydrologic perturbations that result from both climate change and human impact.

Keywords: Ecohydrology; Integrated Water Resources Management; Functional Flows; Ecosystem Services; Hydrological Connectivity; Adaptive Governance; Water Security.

(Submitted: April 02, 2025; Revised: April 18, 2025; Accepted: May 29, 2025; Published: June 30, 2025)

I. Introduction

For most of the 20th centuries, management of water resources was almost solely defined through civil and hydraulic engineering and based on managing and controlling water flows for human benefit (Brauman, 2015; Shen & Chen, 2010; Lenton & Muller, 2012). Such an engineering approach exclusively focuses on mechanical elements like storage, conveyance, and withdrawal of water in support of rapidly growing urban populations, industrial, and agricultural needs (Grizzetti et al., 2016). The riverine landscape of flows was abstracted into static inputs to be manipulated by a system of dams, dikes, and diversion channels; yield maximization and risk of flood avoidance were the prime directives (Jin et al., 2015; Jin et al., 2018; Shaad et al., 2022). Ecological constraints were largely overlooked or included as a minor factor because the management assumption was that flow could be regulated with little collateral impact on aquatic environments.

The results of such segmented management, in fact, account for extensive, even irreversible degradation. Regulation of the riverine flows-that is to say, the magnitude, frequency, and timing of river discharge (riverine flow regime)-effectively dismembered the linkages between rivers, riparian areas, and floodplains. Disconnected systems lead to severely reduced aquatic biodiversity, loss of wetlands, weakened nutrient cycles, and disrupted sediment transport that nourishes healthy rivers (Zalewski, 2015; Liu et al., 2013; Yang et al., 2021). As aquatic connectivity is severed, natural eutrophication processes in water bodies are limited, as is the capacity of aquatic systems to maintain fish populations and recover from human-caused environmental stress (Bunch et al., 2011; Ako et al., 2010). The impact is a weakened capacity of the aquatic

ecosystem to cope with growing pressure from climate change and pollution (Vollmer et al., 2018; Overton et al., 2014; Vogel et al., 2015).

- Synthesize and summarize the state of scientific knowledge on ecohydrological processes and address the shortcomings in current water governance practices.
- Propose a comprehensive framework that incorporates ecological vitality as a central technical design requirement (rather than as a secondary environmental constraint) into water resource management.
- Connect physical hydrology modeling to biological performance metrics, enabling informed decisions about river basin management.
- Develop a set of recommended management interventions that reconcile human water security demands with sustainable ecosystem service provision.

The organization of the paper starts with an introduction of past engineering-driven approaches to water management and their environmental consequences (Section I: Introduction). Then it presents the interactions among hydrological drivers and ecosystem processes (Section II: Conceptual Framework), followed by the compilation of multi-metric models that show the responses of biodiversity to changing flows (Section III: Synthesis). The proposed Integrated Eco-Governance Framework and its foundations are described in Section IV: Proposed Management Framework, accompanied by a table of hydrological drivers and corresponding strategies to build resilience. Finally, conclusions and policy recommendations regarding adaptive governance, legally binding protection of environmental flows, and ecological monitoring embedded in management schemes are discussed in Section V: Conclusion and Policy Recommendations, with references provided.

II. Conceptual Framework

Hydrology and ecology are coupled, at least in principle, because aquatic organisms evolved in pace with the dictates of the hydrological cycle. Physical components such as discharge, baseflow recession, and temperature are not only "physical conditions" or "environmental states," but they are, in the primary sense, ecological signals. The seasonal river pulse, for example, is one of the main sources of aquatic productivity because it is responsible for transporting nutrients between the river channel and the floodplains. Flow is critical for maintaining hydrological cycles: inflow brings organic matter and sediment, which are required for production downstream; outflow keeps diverse habitats, such as the upstream spawning reaches and the downstream nursery reaches, connected.

Biological processes also have a profound effect on the hydrological state of an ecosystem, creating a feedback loop. Vegetation in riparian areas and wetlands acts as a biological regulator for water processes through increasing roughness, thus enhancing recharge and slowing runoff, and enhancing stream bank stability, thus reducing sediment deposition. The rate at which the ecosystem turns over nutrients is also highly influenced by biological processes, as it is the specific microbial populations and the plant community that influence the speed at which nitrogen and phosphorus are metabolized or stored. Healthy biological components of the system increase the water quality and buffering capacity of the ecosystem, which has a significant impact on the "hydrological behavior" of the landscape.

The persistence of an aquatic ecosystem requires a continuous flow of groundwater and surface water. Ground-water discharge provides a crucial "thermal refuge" and base flow during drought or periods of high-water temperatures and acts as a vital base flow, maintaining habitat volume when surface water discharge has diminished to nothing. This interaction has the potential to be lost during management if it ignores the groundwater-surface water connectivity that keeps these populations alive. If it regards hydrology as the "stage" for biology, it will see that alterations to the former will also alter the latter, and a move toward viewing water not as a resource to be managed as a commodity but as a living system is necessary.

III. Synthesis

Current research moves beyond traditional, engineering-focused water management through the use of multi-metric models treating hydrology and biology as coupled parameters. Predictive models such as Environmental Flow Components (EFC) and Ecological Limits of Hydrological Alteration (ELOHA) have been successful in managing systems not on simple low flow thresholds but on recreation of natural flow regimes. It is the functional flows (those flows that create biological events such as spawning for fish, nutrient pulses into floodplains, and flushing of sediments from systems) rather than static flows that maintain aquatic biodiversity. These models utilize georeferenced data combined with biological survey data and successfully predict impacts of certain hydrologic alterations, such as dam-induced flow stabilization, leading to a reduction in habitat diversity and specialists.




Figure 1: Relationship between Functional Flow Regimes and Aquatic Biodiversity Index

Figure 1 shows the effects that three different flow regimes (e.g., natural seasonal pulse, dam-altered flow, and reduced baseflow) can have on aquatic biodiversity indices (species richness, sensitive species abundance, and Shannon diversity index). The x-axis would be the type of flow regime or hydrological modification, and the y-axis would be the value of the biodiversity index. Each line or point could represent a river segment or year of measurement. This would also illustrate the non-linear relationship between hydrological alteration and ecosystem health as seen from the feedback included in the conceptual framework.

Additionally, state-of-the-art hydro-ecological models are starting to include the interaction of groundwater-surface water as another ecosystem stress metric, where modeling with integrated hydro-ecological models has shown a high correlation between baseflow loss-often a result of over-extraction of groundwater-and the alteration of thermal refuges which support cool-water fishes in extreme summer heat. By quantifying the effect that these changes in physical regime can have on biological parameters, scientists have shown that ecological decline often is a non-linear response to crossing hydrological thresholds. This confirmation proves that the integration of ecological indicators into management models would result not only in positive trends in biodiversity but also in an increase in the overall resilience to climate change and hydrology as well.

IV. Proposed Management Framework

To remedy this fragmented nature of water management, it proposes an Integrated Eco-Governance Framework (IEGF) of "Dynamic Ecosystem-Responsive Management." This moves water management from an allocation scheme to an adaptive, responsive, and data-driven regime, which sees the river as an ecosystem as opposed to solely a source of supply. Nested management with ecological flow defined as a "first-order" right, much like water rights for agriculture or municipalities. Would have the ecological pulse needed for biological integrity enshrined in law as opposed to a secondary benefit.

Table 1: Summary of Hydrological Drivers, Ecosystem Functions, and Resilience Implications

Hydrological Driver	Metric/Indicator	Ecosystem Function Impact	Observed Effect	Reference
Flow timing & magnitude	Peak discharge, pulse frequency	Fish spawning, nutrient transport	High natural pulse → increased spawning success	Bunch et al., (2011)
Groundwater baseflow	Baseflow volume	Thermal refugia, habitat continuity	Reduced baseflow → stress in cold-water species	Bunch et al., (2011)
Sediment transport	Suspended solids, sediment load	Habitat formation & substrate quality	Stabilized flow → reduced sediment replenishment	Bunch et al., (2011)
Riparian vegetation	Coverage, density	Bank stability, runoff modulation	Degraded vegetation → increased erosion	Bunch et al., (2011)

Table 1 also explicitly correlates the primary hydrological drivers with the results of ecological impact and measures for resilience, thereby focusing on the association between the use of adaptive management and functional flow regimes to ensure the proper functioning of ecosystems. It also further ties the work to hydrologic resilience in a shifting climate by citing.

Key Framework Pillars

- Environmental Flows (E-Flows) as Dynamic Blueprints: Instead of minimum base flow criteria, this framework requires the use of "functional flow regimes" specific to local species life-cycle needs and nutrient transport requirements, meaning that infrastructure release schedules are proactively managed to simulate natural hydrological events (e.g., Seasonal floods, pulses) which provide the connectivity required for key habitats and induce biological spawning.
- Integrated Stakeholder Negotiation: The IEGF framework utilizes an approach labeled "Benefit Sharing". This technique repositions the dialog between agricultural, industrial, and environmental managers from competition over limited, zero-sum resources to shared and sustained use and management. When water-users are rewarded-for example, through rebates for water efficiency or tradeable permits linked to an indicator of aquatic health-they have an incentive to coordinate water use and schedule operational time in accordance with the rhythms of aquatic systems, thus maintaining access over time.
- Adaptive Feedback Governance: The model applies the Ecohydrological Feedback Loop (see Figure 1), which creates a real-time monitoring and feedback loop that enables water management allocation adjustments to be made based on real-time indicators of ecological health. In this way, water management allocations can remain adaptive and be readily adjusted in real time in response to biological indicators of stress, thereby maintaining the health of the aquatic environment.

V. Conclusion & Policy Recommendations

Bringing together hydrology and ecosystem functioning is a fundamental means of mitigating the impacts of the past engineering-driven water management system. By building and recognizing the two-way interaction between the physical forcing of hydrologic processes and biological feedback mechanisms, management systems can move away from static supply-based models to dynamic models designed to provide resilient ecosystem functioning. Treating water as a fundamental part of the living landscape that provides services such as nutrient processing, flood attenuation, and habitat is made possible by linking the two systems. Implementation of such a management regime begins with governments legally

defining environmental flows from being secondary constraints and becoming the primary, protected right of users by establishing a 'functional flow' regime that emulates natural patterns of variation. Adaptive governance requires managers to build real-time ecological monitoring into their decision-making framework so that managers can flexibly adjust water allocation decisions in direct response to indicators of biological distress. "Benefit-sharing" negotiation systems are required from governments in order to get agriculture, industry, and resource managers to work together through the use of incentives such as efficiency subsidies and tradable water permits directly linked to ecological outcomes. The establishment of multi-metric modeling capacity, recognizing subsurface connectivity and thermal refugia instead of solely water volumes, will be essential for the future long-term security of the resource under a changing climate. Embedding these management practices will allow decision makers to satisfy human water security demands while simultaneously ensuring ecosystem functioning for the future.

References

[1] Zalewski, M. (2015). Ecohydrology and hydrologic engineering: regulation of hydrology-biota interactions for sustainability. *Journal of Hydrologic Engineering*, *20*(1), A4014012. https://doi.org/10.1061/(ASCE)HE.1943-5584.0000999

[2] Vollmer, D., Shaad, K., Souter, N. J., Farrell, T., Dudgeon, D., Sullivan, C. A., ... & Regan, H. M. (2018). Integrating the social, hydrological and ecological dimensions of freshwater health: The Freshwater Health Index. *Science of the Total Environment*, *627*, 304-313. https://doi.org/10.1016/j.scitotenv.2018.01.040

[3] Liu, S., Crossman, N. D., Nolan, M., & Ghirmay, H. (2013). Bringing ecosystem services into integrated water resources management. *Journal of environmental management*, *129*, 92-102. https://doi.org/10.1016/j.jenvman.2013.06.047

[4] Overton, I. C., Smith, D. M., Dalton, J., Barchiesi, S., Acreman, M. C., Stromberg, J. C., & Kirby, J. M. (2014). Implementing environmental flows in integrated water resources management and the ecosystem approach. *Hydrological Sciences Journal*, *59*(3-4), 860-877. https://doi.org/10.1080/02626667.2014.897408

[5] Brauman, K. A. (2015). Hydrologic ecosystem services: linking ecohydrologic processes to human well-being in water research and watershed management. *Wiley Interdisciplinary Reviews: Water*, *2*(4), 345-358. https://doi.org/10.1002/wat2.1081

[6] Yang, D., Yang, Y., & Xia, J. (2021). Hydrological cycle and water resources in a changing world: A review. *Geography and sustainability*, *2*(2), 115-122. https://doi.org/10.1016/j.geosus.2021.05.003

[7] Vogel, R. M., Lall, U., Cai, X., Rajagopalan, B., Weiskel, P. K., Hooper, R. P., & Matalas, N. C. (2015). Hydrology: The interdisciplinary science of water. *Water Resources Research*, *51*(6), 4409-4430. https://doi.org/10.1002/2015wr017049

[8] Bunch, M. J., Morrison, K. E., Parkes, M. W., & Venema, H. D. (2011). Promoting health and well-being by managing for social–ecological resilience: the potential of integrating ecohealth and water resources management approaches. *Ecology and society*, *16*(1), 1-18. https://doi.org/10.5751/es-03803-160106

[9] Jin, G., Wang, P., Zhao, T., Bai, Y., Zhao, C., & Chen, D. (2015). Reviews on land use change induced effects on regional hydrological ecosystem services for integrated water resources management. *Physics and Chemistry of the Earth, parts A/B/C*, *89*, 33-39. https://doi.org/10.1016/j.pce.2015.10.011

[10] Jin, G., Deng, X., Hasan, S. S., Zhao, C., & Gibson, J. (2018). Hydrological ecosystem services for integrated water resources management. *River Basin Management*, 1-27. https://doi.org/10.1007/978-981-10-0841-2_6-1

[11] Shen, Y., & Chen, Y. (2010). Global perspective on hydrology, water balance, and water resources management in arid basins. *Hydrological Processes: An International Journal*, *24*(2), 129-135. https://doi.org/10.1002/hyp.7428

[12] Lenton, R., & Muller, M. (2012). Integrated water resources management in practice, *better water management for development*. Routledge. https://doi.org/10.4324/9781849771740

[13] Shaad, K., Souter, N. J., Vollmer, D., Regan, H. M., & Bezerra, M. O. (2022). Integrating ecosystem services into water resource management: an indicator-based approach. *Environmental Management*, *69*(4), 752-767. https://doi.org/10.1007/s00267-021-01559-7

[14] Grizzetti, B., Lanzanova, D., Liquete, C., Reynaud, A., & Cardoso, A. C. (2016). Assessing water ecosystem services for water resource management. *Environmental Science & Policy*, *61*, 194-203. https://doi.org/10.1016/j.envsci.2016.04.008

[15] Ako, A. A., Eyong, G. E. T., & Nkeng, G. E. (2010). Water resources management and integrated water resources management (IWRM) in Cameroon. *Water Resources Management*, *24*(5), 871-888. https://doi.org/10.1007/s11269-009-9476-4