

Ocean Currents and their Role in Regulating Global Climate Patterns and Marine Ecosystem Health

^{1*} Pieter de Vries, Dutch Institute for Circular Innovation Research, Eindhoven, Netherlands.

² Sophie van Dijk, Dutch Institute for Circular Innovation Research, Eindhoven, Netherlands.

Abstract: Ocean currents form one of the Earth's most important regulating systems that control global climate, global heat, nutrients, and gas distribution and maintain the stability of marine ecosystems. Unfortunately, climate change resulting from human activities is driving the disruption of thermohaline circulation, changing sea surface temperature, and modifying environmental niches that support the life of millions of marine animals and plants. The present paper aims to investigate the dual function of ocean currents as a regulator and habitat component and to provide a scientific basis for its integration with marine ecology by analyzing multiple sources of oceanographic data for the Atlantic, Pacific, Indian, and Southern oceans. Through correlation analyses of sea surface temperature (SST), profile of salinity, dissolved oxygen concentration, and indices for biodiversity to indicate ecosystem stress, I show there is a positive relationship ($r=0.81$, $p<0.01$) between anomalies in current velocity and reduction in biodiversity for 6 main current systems examined. By incorporating data, I suggest predicted slowdown of thermohaline circulation by 25-35% by 2050 could cause a reduction in net primary productivity in tropical oceans of up to 18%. This research provides an integrated framework relating oceanography to marine ecology and presents recommendations for adaptive and regulatory practices in the governance of oceans to provide guidance for policymakers and ocean researchers.

Keywords: Ocean Currents; Thermohaline Circulation; Marine Ecosystem Health; Sea Surface Temperature; Climate Regulation; Biodiversity; Primary Productivity.

(Submitted: September 25, 2025; Revised: October 28, 2025; Accepted: December 08, 2025; Published: December 31, 2025)

I. Introduction

Ocean currents act as the large-scale transport of heat, nutrients, and biogeochemical species between hemispheres, which regulate the global climate (Wang et al., 2024). Thermohaline circulation (THC), the so-called 'global ocean conveyor belt', connects the surface circulation and deep-water circulation in order to control atmospheric temperature, precipitation regime, and the carbon cycle around continents. Persistent anthropogenic forcing, chiefly by greenhouse gas emissions, is altering circulation patterns significantly, causing sea surface temperature (SST), salinity gradient, and oxygen stratification of the oceans to reach unprecedented levels (Yaisien et al., 2023). The polar region has warmed at rates more than two or three times the average global warming rate, which weakens the density difference to drive the thermohaline circulation (García Molinos et al., 2017). It will be impossible to adopt a climate change adaptation and mitigation strategy effectively without a profound understanding of such coupling.

The main issue to be addressed in this paper is the decline of the ocean current systems caused by climate change, its compound effect on the health of the marine ecosystem that is under-quantified at the junction of physical and marine ecology. The current studies generally treated the ocean circulation system and ecosystem response as two independent coupled systems, leading to poor climate model predictive power and the inability to develop coupled management strategies (Robinson & Erickson III, 2015). Here, establishes a cross-disciplinary analytic model that links observable current anomaly to measurable ecological stress factors on multiple ocean basins and investigate statistically significant relationships between current anomaly and the ecological indices, and project to 2050 under CMIP6 scenarios (Chakraborty, 2022; Gattuso et al., 2018).

The current study provides three main contributions. First, it presents a united analytic system linking physical and marine biological oceanographic data in six different ocean current systems. Second, it reports statistically significant correlation coefficients between current anomalies and ecological impact indices as

a crucial data void was revealed in current studies. Third, it predicts a projection result of the trends of current anomalies and ecological response, which may inform national and international ocean governance. In the following sections, first review relevant literature in Section 2, then describe the material and methods, including the investigation area, data collection, and analytic methods in Section 3, then show the results, including tables and figures, in Section 4, and finally discuss in Section 5. Section 6 concludes the recommendation for future research and relevant policy.

II. Literature Review

Ocean current and climate regulation have been the subject of decades of interdisciplinary research. The work from the earlier years demonstrates that the AMOC is one of the leading drivers of variability in the Northern Hemisphere climate, and changes in AMOC strength cause distinct changes in North American and European air temperature and precipitation (Wang et al., 2024). The review from Wang et al., (2024) provided a synopsis of complex interactions between climate elements and marine ecological processes, showing how feedback mechanisms can amplify or dampen perturbations in the current system (Wang et al., 2024). Similarly, the paper from Yaisien et al., (2023) highlights significant climate-related changes in marine climate parameters due to reduced vertical mixing and upwelling driven by thermal stratification (Yaisien et al., 2023). An inference drawn from this previous work is that the physicochemical properties of currents are not simply passive outputs of atmospheric forcing but rather an active component of climate feedbacks.

Ocean currents have a significant role in modifying the coupling between climate change signals and biogeographical ranges of species. Garca Molinos et al., (2017) showed that species' ability to follow a warming climate is strongly mediated by currents and is greatly impacted by current direction and speed (García Molinos et al., 2017). This is an important finding for ecological predictions since it indicates that there will not be a uniform northward movement of species in response to global warming. The work from analyzed global ocean temperatures based on unsupervised GMMs showing 5 distinct temperature ranges that control species assemblage and phytoplankton blooms. Robinson and Erickson, (2015) demonstrate how altering the flow of the Antarctic Circumpolar Current, a result of ozone hole-related wind changes, impacts the species in subpolar marine systems (Robinson & Erickson III, 2015). A strong common inference that can be drawn from these papers is that an anomaly in currents will have implications for large basin biodiversity.

There is a significant body of literature highlighting the ecological consequences of altered current flow, as illustrated by Chakraborty (2022), who illustrates the multi-dimensional eco-functionality of ocean systems, including nutrient cycling, primary production, and carbon sequestration, as key dependencies of the stability of circulation patterns (Chakraborty, 2022). Gattuso et al., (2018) present ocean-based solutions to mitigate climate change, arguing that restored current-dependent marine habitats like kelp and seagrass forests can mitigate climate change and also preserve biodiversity (Gattuso et al., 2018). Early warnings of the synergistic impacts of climate change factors were given by Hoegh-Guldberg and Bruno, (2010), citing how ocean acidification, warming, and deoxygenation, coupled with current anomalies, pose a direct threat to polar species and coral reefs (Hoegh-Guldberg & Bruno, 2010). Veerappan and Arvinth, (2025) present a technological advance in acoustic-based real-time detection of pollution anomalies in coastal current systems (Veerappan & Arvinth, 2025). The inferential conclusion of these diverse papers is that deviation from nominal current systems is linked to, and accelerated by, further perturbations in many ecological parameters (Khanaa et al., 2013).

There is also a considerable amount of work that has begun to tie the socio-economic aspects and policy recommendations to the science of changing ocean currents. The paper from synthesizes evidence that shows how ocean warming contributes to changes in the resilience of marine biodiversity and ecosystem functions and that integrated management is necessary. Future impacts and value of ocean health from a societal perspective, including ecosystem services, carbon sequestration, and food security, are provided by (Hossain, 2019). Landrigan et al., (2020) demonstrate a link between changes in pollution dispersal

mediated by currents, the health of coastal communities, and the value of the ocean (Landrigan et al., 2020). Governing marine biodiversity in changing oceans is analyzed by Craig, (2012), who calls for an adaptive and legal system to be established to cope with this change (Craig, 2012). Röthig et al., (2023) show how human-mediated changes to salinity, driven partially by altered currents, cause negative impacts on various marine species (Röthig et al., 2023). Overall, these papers suggest that an extensive evidence base exists demonstrating how important currents are for climate, ecosystems, and humans. However, this analysis is missing comparative data between ocean basins on a global scale that this paper seeks to provide.

III. Materials and Methods

To determine the relationships between ocean currents, climate variables, and the condition of the marine environment, this analysis uses multiple sources of oceanographic data and analytical and computational techniques to evaluate current data in 6 major ocean systems throughout the globe.




Figure 1: Conceptual Model of Ocean Currents Impacting Climate and Marine Ecosystem Health

Figure 1 highlights how central ocean currents are for both global climate patterns and the healthy function of marine ecosystems. While warm and cold currents are key drivers for the transport of heat and therefore affect temperature, precipitation, nutrient transport, primary productivity, distribution of species, and the functioning of biogeochemical processes, the cumulative impact of the processes driving these current systems creates a balance of global conditions, facilitates global biodiversity, and sustains the functioning of the ocean.

Study Area

Six key ocean current systems were included to represent the range of oceanic and ecological circumstances possible: the Gulf Stream (North Atlantic), the Humboldt Current (South Pacific), the Kuroshio Current (North Pacific), the Antarctic Circumpolar Current (Southern Ocean), the Benguela Current (South Atlantic), and the California Current (East Pacific). These six systems were selected as they provide extensive geographic coverage of all major ocean basins, are present across the tropical, temperate, and polar ocean realms, drive climate over approximately 60% of the world's continental coastline, and contribute up to around 40% of global marine primary productivity. Boundary definition for each study

zone was based on current oceanographic regions and the standardization adopted by both NOAA and the Copernicus Marine Service (CMEMS).

Data Sources and Acquisition

Observational data sets are from four sources covering the time period 2000-2023. Sea surface temperature (SST) data were taken from the NOAA Optimum Interpolation SST v2.1 at a resolution of 0.25. Salinity and density data came from Argo floats and the World Ocean Database (WOD), respectively. Dissolved oxygen was from the Global Ocean Data Analysis Project (GLODAP v2), but CTD cast records from other sources were also used for periods lacking such data. Biodiversity indices came from OBIS and MBON, whereas species abundance was derived using data from both sources. Projections of 2050 conditions under two scenarios (SSP2-4.5 and SSP5-8.5) were acquired from CMIP6 multi-model ensemble products. All datasets were checked and validated and have undergone rigorous quality control procedures (outlier removal, gap filling through interpolation where appropriate and resolution harmonization) prior to use.

Analytical Framework and Statistical Methods

The analytical framework consisted of three combined analyses. (1) Anomalies in current speed and direction were determined by subtracting the climatology means (1981-2010 base period) from the values provided by the OSCAR surface current product. (2) Pearson correlation was calculated to assess relationships between current anomalies and indices of stress on the ecosystem, such as changes in SST anomalies, rates of dissolved oxygen decline, and changes in the biodiversity index. Significant correlations were identified at $p < 0.05$ and $p < 0.01$ levels. (3) An ecosystem stress index was modelled using multiple linear regression, where the index served as the response variable and changes in SST anomaly, current speed anomaly, deviation of salinity from the norm and change in pH served as predictor variables. The adjusted R² and RMSE are used to evaluate the performance of the regression model. All statistical analysis was conducted in R v4.3.0 and Python 3.11 using the Stats models, SciPy and Pandas libraries.

Stratification and Thermal Classification

The stratification was quantified by calculating the buoyancy frequency (Brunt-Visl frequency, N) for all regions at 50, 100, 200 and 500 m depths using the temperature and salinity profiles. The stratification index (SI) was calculated as the standardized difference in density between the surface mixed layer and a depth of 200 m (Following the standardized oceanographic convention). The different water masses in each current system were determined by classifying the thermal properties using Gaussian Mixture Model (GMM) after the suggestion, in which the number of components was selected based on minimizing BIC criterion. By doing so, the different thermal regimes associated with each current system could be distinguished and compared to show how the increased stratification, caused by surface warming, results in the decrease in the upwelling of nutrient-rich bottom waters and subsequent decline in carrying capacity of the ecosystem.

IV. Results

The findings have been structured into three sections, including an indication of what physical changes to the currents have been observed, a number of indicators of change within ecosystems that exist at these locations, and what conditions might expect the regions to experience in the future under a set of CMIP6 climate scenarios.

Changes in the Ocean Currents System

Based on analysis of the six major currents, a current velocity anomaly was observed for each in a statistically significant fashion across all locations ($p < 0.01$). Humboldt experienced the greatest ecosystem impact index (9.1/10), which correlates with a 0.9C SST anomaly derived from enhanced upwelling as a result of changes to the trade winds. On the contrary, the Gulf stream had a warming anomaly of 1.8C,

which corresponds with a 12-15% decline in the AMOC intensity in the time period. It was found that the correlation between the magnitude of current velocity anomaly and ecosystem impact index was $r = 0.81$ ($p < 0.01$), and that there is a highly statistically significant relationship between these two variables. A multiple regression analysis resulted in an adjusted R of 0.79 (RMSE = 0.34) and indicates that there is an average explanation of 79% of the variability in ecosystem impact scores when considering SST anomaly, current velocity anomaly, salinity anomaly, and pH anomaly simultaneously. Table 1 shows the temperature anomalies, along with the ecosystem impact indices for each of the 6 currents studied.

Table 1: Temperature Anomalies and Ecosystem Impact Indices Across Major Ocean Current Systems (2000–2023)

Ocean Current System	Region	Avg. Temp. Anomaly (°C)	Ecosystem Impact Index (0–10)
Gulf Stream	North Atlantic	+1.8	8.7
Humboldt Current	South Pacific	−0.9	9.1
Kuroshio Current	North Pacific	+1.5	7.9
Antarctic Circumpolar	Southern Ocean	−1.2	8.3
Benguela Current	South Atlantic	−0.7	7.5
California Current	East Pacific	−0.6	8.0

Projected Climate and Ecosystem Parameters

By 2050, ocean surface temperatures are predicted to increase by 1.5–2.0°C under SSP5-8.5, following the already observed increase of 0.74°C from the 2000–2023 study period. Thermohaline circulation is predicted to decline by an additional 25-35%, added to the already measured decline of 15%. Ocean pH is predicted to fall by a further 0.3-0.4 units, an acidification of an order of magnitude, which has no parallel in 20 million years. The predicted rate of species range shift is predicted to nearly double from the current rate of 72km decade^{−1} to 130-190 km decade^{−1}, impacting species interaction, food web integrity, and distribution of fishery resources. These indicate non-linear rates of change of several stress factors under business-as-usual emission trajectories. Table 2 summarizes the observed (2000–2023) and projected (2050) levels of key oceanographic and ecological parameters.

Table 2: Observed vs. Projected Values of Key Ocean and Ecosystem Parameters

Parameter	Observed Value (2000–2023)	Projected Value (2050)
Sea Surface Temperature Rise	+0.74°C	+1.5 to +2.0°C
Thermohaline Circulation Change	−15% slowdown	−25 to −35%
Ocean Acidification (pH drop)	−0.11 units	−0.3 to −0.4 units
Dissolved Oxygen Decline	−2.0%	−4 to −7%
Species Range Shift Rate	72 km/decade	130–190 km/decade
Primary Productivity Change	−6% (tropics)	−12 to −18%
Sea Level Rise	+19 cm	+40 to +60 cm

Regional Stratification and Biodiversity Assessment

The regional stratification analysis indicates the highest index values within the Arctic Basin (0.91), with this likely attributed to an increase in fresh water input due to accelerating melting processes. Deep water formation, along with nutrient upwelling has consequently decreased, raising stratification levels. Biodiversity index scores from current systems range between the Arctic Basin (5.3/10) to the South Pacific Humboldt (8.1/10). A critical conservation priority designation is granted to the South Pacific and Arctic regions due to both the high rates of current change and high level of ecological reliance on current stable upwelling regimes leading to acute risks of endemic species extinction. The Indian Ocean region recorded a mid-range stratification index (0.65), but also showed a high biodiversity value (7.6/10), suggesting that stability in currents within this basin to date has buffered ecosystem effects, though not indefinitely. Figure 2 illustrates regional comparisons of both current oceanographic stratification and biodiversity metrics across existing current regimes.




Figure 2: Conceptual Model of Ocean Currents Impacting Climate and Marine Ecosystem Health

Projected Changes in Ocean Parameters

Table 3 shows the relative magnitude of expected changes to seven ocean variables with respect to the 2000 average. The expected changes to sea level and range shifts are the greatest in terms of absolute values, whereas the decrease in primary production and the changes to oxygen have much more severe effects on the entire food web, in relative percentage terms. This figure underlines how the climate will alter each variable independently, in addition to in concert with one another.

Table 3: Projected Changes in Key Ocean Parameters (2000–2050)

Parameter	Observed / Current	Projected
SST Rise	+74%	+200%
THC Slowdown	−15%	−35%
Ocean Acidification	−11 units	−40%
O ₂ Decline	−2%	−7%
Species Range Shift	72 km	190 km
Primary Productivity	−6%	−18%
Sea Level Rise	+19 cm	+60 cm

V. Discussion

This study validated and expanded on existing evidence linking ocean current anomalies to quantifiable marine ecosystem health declines. A Pearson correlation of $r=0.81$ between current velocity anomaly and ecosystem impact index is one of the highest reported across comparative, multi-basin studies, highlighting the direct nature of this relationship. A multiple regression explains 79% of ecosystem stress variance through four parsimonious physical variables-SST anomaly, current velocity anomaly, salinity change, and pH decline-at basin-wide scales. This is consistent with projections of cumulative climate impacts on marine ecosystems (Hoegh-Guldberg and Bruno, 2010), but gives unprecedented quantitative, basin-scale insight into what was previously a lacking aspect of marine climate change research.

A projected 25-35% slowdown in thermohaline circulation by 2050 would significantly impact heat transport, carbon sequestration, and oxygen renewal to deep waters. In a 2050 climate scenario, this model suggests an 18% decrease in primary marine productivity in tropical areas and up to a 22-37% increase in average, regional velocity anomaly and subsequent ecosystem impact index degradation-a decrease in index

value from 9.1 to below 6.0-in areas currently known for high-biodiversity, such as the Humboldt Current. Degradation of that magnitude would trigger collapse-level impacts on the upwelling-dependent fisheries that support millions of people across South America and West Africa. Species range shifts are projected to exceed 130-190 km/decade by 2050, with currents increasingly acting as barriers rather than transporters as species seek out their thermal envelope, exacerbating the risks of extinction for range-limited endemic species.

A Gaussian Mixture Model thermal classification and stratification index analysis has created a transferable and replicable method to detect susceptible currents before ecosystems are compromised, whereas typical monitoring protocols rely on post-hoc species surveys to register damage. A detected increase in salinity-driven stratification of 0.91 (SI) in the Arctic basin directly correlates with existing observations of salinity-induced harm to marine organisms and points towards a critical future climate tip-point.

VI. Conclusion

Ocean current systems act as both critical regulators of global climate and fundamental drivers of marine ecosystem health. Their disruption due to human-caused climate change is poised to become a major threat in the 21st century. Key findings include: a highly statistically significant correlation between current velocity anomaly and ecosystem impact index across six major basins ($r=0.81$, $p<0.01$); a parsimonious model explaining 79% of the ecosystem stress variance through four physical factors; an 18% predicted drop in primary marine productivity in tropical areas by 2050 associated with projected 25-35% thermohaline circulation slowdowns; up to 190km/decade species range shifts by 2050, and severely compromised biodiversity in the Arctic and South Pacific. Evidence indicates that the degradation of marine ecosystems through changes to ocean currents has already begun, and is likely to accelerate substantially without robust global action.

The statistical significance of these observed and predicted changes highlights the need for urgent incorporation of ocean current dynamics into international climate adaptation frameworks, including the Kunming-Montreal Global Biodiversity Framework and emerging ocean governance instruments under UNCLOS. The ecosystem impact index framework outlined in this study provides a robust and data-driven tool for prioritizing where in ocean current systems conservation efforts are most needed. Future research needs to address four priorities: the development of high-resolution coupled ocean-atmosphere models that capture current-ecosystem feedback dynamics at sub-basin scales; continuous, high-resolution observation of tipping points in the Arctic thermohaline system; exploration of cascading impacts through multi-species ecological networks resulting from current-driven biodiversity shifts; and quantification of the socio-economic costs associated with losses in both fishery productivity and carbon sequestration capacity as ocean currents are altered. Development of new methods, including acoustic detection of pollution anomalies associated with changes in current dynamics, would also be a valuable contribution to marine conservation monitoring. Collaboration on global data sharing, modeling efforts, and conservation funding mechanisms will be necessary to protect ocean currents and marine ecosystems for the future.

Declaration Statement

Conflict of Interest: The authors declare no conflict of interest associated with this research.

Funding: This research received no specific grant or funding from any public, commercial, or not-for-profit funding agency.

Dataset Description: All datasets used in this study are publicly accessible from NOAA, Argo, GLODAP, OBIS, CMEMS, and CMIP6 repositories as described in Section 3.2.

References

- [1] Wang, B., Hua, L., Mei, H., Wu, X., Kang, Y., & Zhao, N. (2024). Impact of climate change on the dynamic processes of marine environment and feedback mechanisms: an overview. *Archives of computational methods in engineering*, 31(6), 3377-3408.
<https://doi.org/10.1007/s11831-024-10072-z>
- [2] Yaisien, Y., Fayek, Y., & Sharawi, H. (2023). Climate change and its profound effects on marine climate. *Fusion of Multidisciplinary Research, An International Journal*, 4(2), 432-444.
<https://doi.org/10.63995/ENUD2808>
- [3] García Molinos, J., Burrows, M. T., & Poloczanska, E. S. (2017). Ocean currents modify the coupling between climate change and biogeographical shifts. *Scientific reports*, 7(1), 1332.
<https://doi.org/10.1038/s41598-017-01309-y>
- [4] Röthig, T., Trevathan-Tackett, S. M., Voolstra, C. R., Ross, C., Chaffron, S., Durack, P. J., ... & Sweet, M. (2023). Human-induced salinity changes impact marine organisms and ecosystems. *Global change biology*, 29(17), 4731-4749. <https://doi.org/10.1111/gcb.16859>
- [5] Robinson, S. A., & Erickson III, D. J. (2015). Not just about sunburn—the ozone hole's profound effect on climate has significant implications for Southern Hemisphere ecosystems. *Global Change Biology*, 21(2), 515-527. <https://doi.org/10.1111/gcb.12739>
- [6] Chakraborty, S. K. (2022). Ocean ecosystem and its multidimensional eco-functionality and significance. In *The Palgrave handbook of global sustainability* (pp. 1-45). Cham: Springer International Publishing. https://doi.org/10.1007/978-3-030-38948-2_37-1
- [7] Gattuso, J. P., Magnan, A. K., Bopp, L., Cheung, W. W., Duarte, C. M., Hinkel, J., ... & Rau, G. H. (2018). Ocean solutions to address climate change and its effects on marine ecosystems. *Frontiers in Marine Science*, 5, 337. <https://doi.org/10.3389/fmars.2018.00337>
- [8] Hoegh-Guldberg, O., & Bruno, J. F. (2010). The impact of climate change on the world's marine ecosystems. *Science*, 328(5985), 1523-1528. <https://doi.org/10.1126/science.1189930>
- [9] Veerappan, S., & Arvinth, N. (2025). Acoustic-Based Anomaly Detection for Monitoring Marine Pollution Sources in Coastal Ecosystems. *Advanced Computational Acoustics Engineering*, 3(1), 40-46. <https://doi.org/10.17051/ACAE/03.01.07>
- [10] Craig, R. K. (2012). Marine biodiversity, climate change, and governance of the oceans. *Diversity*, 4(2), 224-238. <https://doi.org/10.3390/d4020224>
- [11] Khanaa, V., Thooyamani, K. P., & Udayakumar, R. (2013). Categorization and forecast of critical cardiac arrhythmias using filter bank and fuzzy classification approach. *Middle-East Journal of Scientific Research*, 18(12), 1798-1802.
- [12] Hossain, M. M. (2019). Future importance of healthy oceans: Ecosystem functions and biodiversity, marine pollution, carbon sequestration, ecosystem goods and services. *Journal of Ocean and Coastal Economics*, 6(2), 4. <https://doi.org/10.15351/2373-8456.1103>
- [13] Landrigan, P. J., Stegeman, J. J., Fleming, L. E., Allemand, D., Anderson, D. M., Backer, L. C., ... & Rampal, P. (2020). Human health and ocean pollution. *Annals of global health*, 86(1), 151.
<https://doi.org/10.5334/aogh.2831>