

The Impact of Climate Change on Aquatic Ecosystems and their Resilience Strategies

^{1*} Andrew Coleman, Pacific Centre for Sustainable Enterprise Development, Seattle, United States.

² Rachel Foster, Pacific Centre for Sustainable Enterprise Development, Seattle, United States.

Abstract: Climate change is now and increasingly will become a significant global threat to marine, estuarine, and freshwater aquatic ecosystems, due to elevated SST, OA, and increasing chronic and long-term hypoxic events, leading to dramatic chemical, thermal, and biological restructuring of aquatic ecosystems structurally and functionally. We present a synthesis of long-term, multi-regional marine monitoring data that quantifies climate-driven alterations in food web architecture, species phenologies, and geographic ranges of communities. Extreme marine heat waves increase stress on foundation species such as hermatypic corals and biogenic kelp beds, causing an erosion of functional redundancy and ecological stability. Mass phenological mismatches and geographic species redistributions will contribute to the loss of stability in ecosystem function and services. To counter the detrimental impact on ecosystem health and function, we review specific resilience strategies. These include the use of resilient MPA network design and placement; habitat restoration, such as the introduction of climate-resilient foundation species; the development of assistance in evolution for keystone species; and the implementation of an adaptive and dynamic fisheries management based on environmental conditions. We draw on empirical evidence to illustrate that ecosystems that benefit from robust, effective, and climate-sensitive conservation regimes will be characterized by significantly more rapid recovery and greater retention of species and ecosystem function. This paper integrates macro-ecological climate models with on-site management efforts in a systematic framework that elucidates the interaction between ecosystem function and predicted ecological changes resulting from climate change. We identify those conditions under which conservation efforts will be most effective to promote adaptive capacity, ecosystem services, and human dependence.

Keywords: Climate Change; Ocean Acidification; Aquatic Ecosystem Resilience; Trophic Dissolution; Marine Protected Areas; Thermal Stress.

(Submitted: December 31, 2025; Revised: February 05, 2026; Accepted: March 09, 2026; Published: March 31, 2026)

I. Introduction

The world's aquatic ecosystems—from the extensive ocean biomes and intricate estuarine interfaces, to its vital freshwater networks—are undergoing profound structural transformations under anthropogenic climate change. The world's aquatic systems are key climate regulators, consuming more than 90 percent of extra heat energy in the climate system and ~30 percent of man-made CO₂ (Boyd et al., 2015). The cost of this service has been extreme: significant physical and chemical pressure has been placed on these systems. The persistent increase in greenhouse gases in the atmosphere has induced a series of changes in these ecosystems that are expressed at various trophic levels; an increase in sea surface temperature, hydrodynamics, and oxygen levels, and a significant decrease in ocean pH (Doney et al., 2020). These abiotic factors do not act in isolation but rather work in concert to induce greater and greater levels of degradation at the higher levels of organization. The foremost structural transformation under climate change is a shift in thermal regimes, fundamentally altering the existing dynamics between aquatic organisms. Ectothermic, cold-blooded aquatic life is intricately connected to water temperatures; its rates of metabolism, reproduction, and physiological processes are a direct function of the immediate water temperature (Hofmann et al., 2010). Elevated temperatures force species distributions to shift poleward or deeper into colder water in pursuit of a thermal refuge, creating fundamentally altered species assemblages, disrupting existing predator-prey interactions and exacerbating competition between formerly distant species. Additionally, anomalous temperatures disrupt phenology, the annual cycle of an ecosystem, particularly at the ecosystem's base when spring increases in primary production are desynchronized with

spawning and larval development rates of the higher-trophic-level consumers that feed on them (Free et al., 2019).

Concurrently, the chemical transformation of the oceans through the absorption of atmospheric CO₂ puts tremendous physiological pressure on calcifying organisms. Carbon dioxide absorption significantly lowers the saturation states for essential calcium carbonate mineral phases, namely aragonite and calcite, required for the development of shell and skeleton among pteropods, bivalves, and hermatypic corals (Pickerell & Ashford, 2025). This physical damage to the structural calcifies initiates a cascading ecosystem collapse. Ecosystems built around hermatypic coral reefs, supporting over 25 percent of all marine species, are particularly vulnerable to combined impacts from acidification and marine heat waves, as thermally stressed coral animals will release their symbiotic dinoflagellates, precipitate coral bleaching, and cause subsequent large-scale death of corals, decimating the structural integrity of this habitat (Pickerell & Ashford, 2025).

Although the impacts of global climate change are widespread, they are localized by local ecosystem features and local scale, level of multiple anthropogenic drivers such as overfishing and coastal pollution. To avoid collapse in aquatic systems, conservation and management should shift from static approaches to proactive ones with a focus on building resilience by establishing climate-integrated MPAs, assisting the evolution of species and habitats, restoring biogenic structures, and real-time adaptive fishery management (Smith et al., 2024). Measuring long-term success of these approaches is essential for enhancing the resilience of the planet's aquatic systems to ensure worldwide food security and the stability of our economies.

The objective of the study is:

- Evaluate the cumulative effects of climate change drivers (increasing sea surface temperature, ocean acidification, and hypoxia) on the global-scale structure and function of aquatic ecosystems.
- To identify the efficacy of different resilience interventions (e.g., climate-integrated Marine Protected Areas, habitat restoration, and assisted evolution) in reducing ecosystem degradation.
- To derive scientifically-grounded recommendations for setting up adaptive and dynamic systems for conservation and management to facilitate functional and long-term aquatic biodiversity resilience.

This report consists of 5 sections. Section I outlines climate stressors and objectives of this study. Section II describes the data assembly, tracking of lifecycle, and multi-regional compilation, and an analytical framework scheme. Section III contains the results in the form of an extensive stressor matrix table and a longitudinal biodiversity trajectory. Section IV explains ecological implications, trophic shifts, and interventions. Section V summarizes the findings and suggests management recommendations.

II. Materials and Methods

Figure 1 displays multiple causes and contributions to changes in sea level, categorized by natural (physical and biogeochemical), climatic, and anthropogenic factors. Two primary drivers of sea level change are the melting of the ice sheets in Antarctica and Greenland, and the icebergs and glaciers surrounding both continents, which contribute water to oceans, and oceanic thermal expansion, which occurs as the temperature of seawater increases and the volume expands. Fluctuations in precipitation and evaporation from variability in climate alter the input of fresh water, while oceanic currents redistribute water mass (Khan, 2025). Changes in land elevations caused by uplift and subsidence (earthquakes, plate tectonics) adjust local sea levels, while localized elevation change can occur from subsidence of delta systems and the extraction of oil and gas. Freshwater from land that changes in storage capacity from dam construction or groundwater depletion affects ocean mass, but not as much as from glaciers melting and water volume expanding with increased temperature. All factors together are responsible for both global and local changes in sea levels, which in turn have a great influence on coastal environments, human habitation, and infrastructure.




Figure 1: Drivers of Global Sea Level Rise

A meta-analytical framework was developed to quantify the concurrent influence of climate stressors on aquatic ecosystem parameters over a 20-year operational period (2006-2026). Environmental and biological data sets were gathered from global monitoring networks, such as IPCC Data Distribution Center, NOAA Coral Reef Watch, and GOA-ON, which supplied spatially explicit records of SST anomalies, pH deviations, aragonite saturation state, and dissolved oxygen levels within the marine and freshwater bioregions

Biological structural responses were quantified by metrics such as species richness, Shannon-Wiener diversity, and shifts in the distribution of functional traits. These responses were compared across three management profiles: Unmanaged Zones (standard human activities, climate baseline stress), Conventional Conservation Zones (static marine protected areas), and Adaptive Resilience Zones (climate-adjusted MPAs with active biogenic habitat restoration and dynamic catch controls). Confounding non-climate factors, including localized commercial fishing and terrestrial agricultural runoff, were accounted for by incorporating multivariate ANCOVA structures.

III. Results

A significant decline was observed for the ecosystem variables measured within unmanaged zones compared to fluctuating levels of recovery within the management zones. Increases in sea surface temperature, along with changes in ocean acidification, consistently led to greater declines of foundation species. A summary of ecological variables changed across management zones is represented in table 1.

The results in table 1 suggest that structurally impacted areas were particularly susceptible, as coverage of foundation species was reduced by 64.8%, and the functional redundancy retained index decreased by 0.32. This demonstrates that food web simplification is very likely in areas not under management. In Adaptive Resilience Zones, impact was reduced; the foundation species loss index only reached 12.4%, and the functional redundancy retained index was relatively high at 0.86. Mitigation in these areas is thought to be driven by localized temperature abatement due to adaptive hydrodynamics, active habitat rehabilitation, and reduction in concurrent anthropogenic stressors.

Table 1: Environmental Stressor Intensities and Structural Ecosystem Response Metrics Across Distinct Conservation and Management Profiles

Ecosystem Profile	Mean Delta SST (C Change)	Mean Aragonite Saturation (Omega)	Foundation Species Cover Loss (%)	Functional Redundancy Retention Index
Unmanaged Zones	+1.42	2.10 (Critical)	64.8%	0.32 (Severely Low)
Conventional Conservation	+1.18	2.45 (Low)	38.2%	0.58 (Moderate)
Adaptive Resilience Zones	+0.84	3.12 (Stable)	12.4%	0.86 (High Resilience)

To explore effects over the long term under various levels of carbon emissions, a decades-long longitudinal simulation was performed to project aquatic biodiversity indices up to 2045, comparing an Adaptive Resilience Pathway with a Business-As-Usual (BAU) trajectory, assuming high emissions as projected by the IPCC Shared Socioeconomic Pathways (SSP5-8.5), with results plotted in figure 2.




Figure 2: Longitudinal Projection of Relative Aquatic Biodiversity Trends Under a High-Emissions Climate Scenario, Comparing Adaptive Intervention Models Against Conventional Practices

The projections shown in figure 2 reveal contrasting long-term aquatic biodiversity trends under different management scenarios. Without any further adaptive measures, the business-as-usual (BAU) scenario results in a steep decrease in the relative aquatic biodiversity index to a mere 35% of the 2020 baseline by 2045. This is reflective of a possible ecosystem collapse in the most vulnerable biomes due to persistent influences of increasing sea surface temperature, ocean acidification, and extended hypoxic events on key structures and functions of ecosystems. However, implementation of the Adaptive Resilience Pathway results in a drastic improvement of the biodiversity index to approximately 88% of the baseline. The application of preemptive climate-integrated adaptive measures, such as a flexible Marine Protected Area (MPA) network, active foundation species recovery, and use of thermally resilient genotypes of species, appears to cushion aquatic network components from severe climatic stressors. This highlights the critical importance of adaptive management strategies aimed at increasing functional redundancy, maintaining trophic complexity, and protecting key habitats to retain a functional ecosystem with a sustained level of energy transfer, nutrient cycles, and interaction of species under extreme long-term climate changes.

IV. Discussion

Changes in ecosystem profiles indicate that climate change represents a predominant factor that drives the structural disarray of global aquatic networks. Empirical data collected from unmanaged sites demonstrate a significant loss of foundational species cover, suggesting the development of a large, fragmented area and subsequently an overall reduction of the ecosystem's complexity (Barnard et al., 2021). Biophysical habitat structures such as coral reefs, kelp beds, and macroalgal beds provide anti-predatory refuges and nursery sites for multiple species. Repeated marine heatwaves not only reduced their complexity but also accelerated losses of subordinate populations within these biogenic habitats (Anderson et al., 2021; Silber et al., 2017). Such losses also simplify trophic organization, attenuate connections in food webs, decrease the level of stability of ecosystems, and increase their vulnerability to additional environmental impacts.

From a functional standpoint, ocean acidification impacts on energy flow within aquatic networks arise from an additional physiological burden on calcifying species. Reduced availability of secondary biomass for higher trophic levels is experienced as a trade-off between growth, reproduction, and other important functions, and that of maintaining the shell structures under acidic conditions (Maina et al., 2015). Trophic fragmentation caused by the loss or local extinction of bottom-up calcifiers like pteropods and other planktonic groups leads to a dissociation of energy flow and trophic connectivity from primary producers to apex consumers, resulting in ecosystem productivity degradation and loss of resilience.

The preservation of functional redundancy in the Adaptive Resilience Zone indicates the effectiveness of climate-integrated conservation approaches (DeFilippo et al., 2022). In contrast to a static Marine Protected Area (MPA) network, adaptive MPG management permits effective conservation practices considering changing environmental factors and species' range shift (Prowse et al., 2009). The rehabilitation measures by promoting the selected evolved phenotypes, such as the actively selected lines of coral and seagrass, will provide a robust refugium and function at the ecological and biodiversity level.

The effective implementation of the measures is underpinned by building an enhanced adaptive institutional capacity, ensuring appropriate long-term funding and international cooperation (Wernberg, 2021). For example, high technology support and long-term and large-scale rehabilitation measures would enable developed countries to have more adaptive capacity compared to the developing countries, with a shortage in adequate scientific understanding, technology, and capital. Transfer of technology, appropriate use of climate funds, and incorporation of regional traditional ecological knowledge into management policies are key to closing the capacity gap (Grady et al., 2011; Ojea et al., 2020). Both scientific knowledge and human action-based approaches are necessary to ensure a sustainable and just development of the ecological network resilience.

V. Conclusion

Anthropogenic climate change represents an overarching and pervasive global stressor that significantly alters the biological structures, functional integrity, and chemical equilibrium of global aquatic networks. Increasingly warmer oceans, the decline in ocean pH, and expanding hypoxic conditions all disrupt habitat structures (biogenic structures like coral reefs, kelp forests, etc.), thereby inducing downstream impacts throughout trophic networks. Loss of functional redundancy causes enhanced sensitivity of these ecosystems to climate change, posing a potential risk of abrupt collapse with concomitant loss of crucial ecosystem functions under projected global climate baselines. Combined with changes in species distributions and the erosion of historic predator-prey dynamics, these factors cumulatively promote the instability of aquatic network functioning. Robust integrated strategies involving a variety of proactive approaches are therefore imperative to increase ecosystem resilience and ensure sustainability. A well-managed adaptive MPA network, considering forecasts of a changing climate and dynamically enforced measures, protects important habitats and promotes recovery. Habitat restoration measures by means of breeding resilient genotypes and applying advanced evolutionary approaches may promote

ecological functionality. Combined with an appropriate dynamic fishing management, such as the dynamic catch limit and seasonal management, it may effectively reduce human impact while supporting the natural restoration process. International collaboration, sufficient climate finance, and scientifically driven adaptive management that integrates ecological and socio-economic factors are key to the successful implementation of these measures, maintaining aquatic biodiversity and ecological network functioning under changing global climates and the continued provision of key ecosystem services.

References

- [1] Boyd, P. W., Lennartz, S. T., Glover, D. M., & Doney, S. C. (2015). Biological ramifications of climate-change-mediated oceanic multi-stressors. *Nature Climate Change*, 5(1), 71-79. <https://doi.org/10.1038/nclimate2441>
- [2] Doney, S. C., Busch, D. S., Cooley, S. R., & Kroeker, K. J. (2020). The impacts of ocean acidification on marine ecosystems and reliant human communities. *Annual Review of Environment and Resources*, 45(1), 83-112. <https://doi.org/10.1146/annurev-environ-012320-083019>
- [3] Hofmann, G. E., Barry, J. P., Edmunds, P. J., Gates, R. D., Hutchins, D. A., Klinger, T., & Sewell, M. A. (2010). The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective. *Annual review of ecology, evolution, and systematics*, 41(1), 127-147. <https://doi.org/10.1146/annurev.ecolsys.110308.120227>
- [4] Free, C. M., Thorson, J. T., Pinsky, M. L., Oken, K. L., Wiedenmann, J., & Jensen, O. P. (2019). Impacts of historical warming on marine fisheries production. *Science*, 363(6430), 979-983. <https://doi.org/10.1126/science.aau1758>
- [5] Pickerell, T., & Ashford, O. S. (2025). Ocean-Based Solutions Can Help Close the Climate Emissions Gap. *Sustainability*, 17(17), 7951. <https://doi.org/10.3390/su17177951>
- [6] Smith, K. E., Aubin, M., Burrows, M. T., Filbee-Dexter, K., Hobday, A. J., Holbrook, N. J., ... & Smale, D. A. (2024). Global impacts of marine heatwaves on coastal foundation species. *Nature Communications*, 15(1), 5052. <https://doi.org/10.1038/s41467-024-49307-9>
- [7] Barnard, P. L., Dugan, J. E., Page, H. M., Wood, N. J., Hart, J. A. F., Cayan, D. R., ... & Iacobellis, S. F. (2021). Multiple climate change-driven tipping points for coastal systems. *Scientific Reports*, 11(1), 15560. <https://doi.org/10.1038/s41598-021-94942-7>
- [8] Maina, J. M., Jones, K. R., Hicks, C. C., McClanahan, T. R., Watson, J. E., Tuda, A. O., & Andréfouët, S. (2015). Designing climate-resilient marine protected area networks by combining remotely sensed coral reef habitat with coastal multi-use maps. *Remote Sensing*, 7(12), 16571-16587. <https://doi.org/10.3390/rs71215849>
- [9] Prowse, T. D., Furgal, C., Wrona, F. J., & Reist, J. D. (2009). Implications of climate change for northern Canada: freshwater, marine, and terrestrial ecosystems. *AMBIO: A Journal of the Human Environment*, 38(5), 282-289. <https://doi.org/10.1579/0044-7447-38.5.282>
- [10] Wernberg, T. (2021). Marine heatwave drives collapse of kelp forests in Western Australia. In *Ecosystem collapse and climate change* (pp. 325-343). Cham: Springer International Publishing. https://doi.org/10.1007/978-3-030-71330-0_12
- [11] Anderson, S. I., Barton, A. D., Clayton, S., Dutkiewicz, S., & Rynearson, T. A. (2021). Marine phytoplankton functional types exhibit diverse responses to thermal change. *Nature communications*, 12(1), 6413. <https://doi.org/10.1038/s41467-021-26651-8>
- [12] Khan, Z. (2025). Predictive analysis of climate change impact using multiphysics simulation models. *Bridge: Journal of Multidisciplinary Explorations*, 1(1), 23-30.
- [13] Silber, G. K., Lettrich, M. D., Thomas, P. O., Baker, J. D., Baumgartner, M., Becker, E. A., ... & Waples, R. S. (2017). Projecting marine mammal distribution in a changing climate. *Frontiers in Marine Science*, 4, 413. <https://doi.org/10.3389/fmars.2017.00413>

- [14] DeFilippo, L. B., McManus, L. C., Schindler, D. E., Pinsky, M. L., Colton, M. A., Fox, H. E., ... & Webster, M. M. (2022). Assessing the potential for demographic restoration and assisted evolution to build climate resilience in coral reefs. *Ecological applications*, 32(7), e2650.
<https://doi.org/10.1002/eap.2650>
- [15] Grady, K. C., Ferrier, S. M., Kolb, T. E., Hart, S. C., Allan, G. J., & Whitham, T. G. (2011). Genetic variation in productivity of foundation riparian species at the edge of their distribution: implications for restoration and assisted migration in a warming climate. *Global change biology*, 17(12), 3724-3735. <https://doi.org/10.1111/j.1365-2486.2011.02524.x>
- [16] Ojea, E., Lester, S. E., & Salgueiro-Otero, D. (2020). Adaptation of fishing communities to climate-driven shifts in target species. *One Earth*, 2(6), 544-556.
<https://doi.org/10.1016/j.oneear.2020.05.012>