

Human Activities and their Impact on Marine Habitat Destruction and Coastal Ecosystems

^{1*} Felipe Rodrigues, Atlantic Brazil Development Innovation Centre, Recife, Brazil.

² Camila Ferreira, Atlantic Brazil Development Innovation Centre, Recife, Brazil.

Abstract: Marine and coastal ecosystems are currently undergoing unprecedented degradation at all spatio-temporal scales under a highly interacting suite of anthropogenic activities. The combined effect of coastal artificialization, industrial pollution, and destructive fishing practices was investigated using data from thirty-eight marine sampling matrices over a time scale and compared against indices of physicochemical properties, benthic structure integrity, and local species diversity, mapped against local human population density and coastal artificial surface index. Empirical analysis shows non-linear declines in coral cover and seagrass density whenever there is an anthropogenic index exceeding 18.2% for the coastal artificial surface index within a 5 km zone, and commercial bottom-trawling strongly correlates with a total loss of benthic structural complexity and macrofaunal functional richness. Analysis by linear mixed-effects modeling shows that the implementation of continuous MPAs and active habitat rehabilitation initiatives significantly buffers marine species from terrestrial sources of pollution. This work provides the first empirically robust, quantitatively explicit decision-making framework for coastal ecosystem managers, suggesting severe local zoning rules to mitigate irreversible ecosystem decline.

Keywords: Marine Anthropocene; Habitat Fragmentation; Coastal Armoring; Coral Reef Degradation; Benthic Complexity; Marine Protected Areas.

(Submitted: March 14, 2026; Revised: April 07, 2026; Accepted: May 22, 2026; Published: June 30, 2026)

I. Introduction

The terrestrial-marine boundary is one of the most biogeochemically active and ecologically sensitive boundaries on the planet. Coastal ecosystems (i.e., salt marsh, mangrove, seagrass, shallow water coral reef systems) supply some of the most critical ecosystem services we rely on, including storm surge buffering, carbon sequestration from the atmosphere, and nursery habitat for economically important world fisheries. Despite their enormous ecological and economic value, these systems are experiencing a period of extreme structural collapse due to over-humanization. The increasing trends of urbanization, industrial development, globalized maritime traffic, and overexploitation have significantly perturbed the stable equilibrium dynamics within these marine environments, and the world's diverse marine communities are moving towards a functional collapse.

The processes by which humans alter the coastal environment are varied and highly consistent across spatial scales. The coastal alteration process for urban development primarily includes land-filling and the construction of fixed structures, or rigid concrete seawalls, port structures, etc. In this manner, the entire localized intertidal and subtidal habitat is eliminated, and regional hydrodynamics and sediment dynamics are altered by replacement with a permanently immobile substrate. In addition, dense urbanized and agricultural areas deliver enormous loads of industrial pollutants (including heavy metals and toxin compounds), plastic macro-debris, and macronutrients to shallow coastal waters from land-derived sources. The continuous input of toxins and eutrophication leads to a chronic toxic environment, which decreases local organisms' physiological resilience (especially foundation species, like corals or marine angiosperms).

Land-based stressors interact strongly with direct anthropogenic impacts on marine environments. Commercial bottom trawling, a highly damaging mechanical disturbance that involves the dragging of heavy gear across the seafloor, literally flattens the complex benthic structure, destroying slow-growing biogenic structures such as sponge field communities and soft coral communities. In this manner, the structure is severely depleted, and fine sediments are resuspended, which can inundate the local environment and destroy complex trophic communities through both smothering and habitat destruction.

This loss of structure decreases macro-faunal biodiversity and changes complex, high-trophic environments to simplistic, low-functioning systems. We aimed to identify threshold values that mark a fundamental shift between systems that can tolerate human disturbances and systems that have been irrevocably degraded by human activity.

While there is a general correlation between human population density and habitat loss, determining the critical ecological threshold at which a marine community collapses remains a difficult question. The majority of current management strategies incorporate spatially broad data sets, which miss important site-specific land use changes and synergistic effects of land-based pollution and bottom trawling. As such, field-based monitoring with high temporal and spatial resolution across clearly defined human disturbance gradients is necessary to isolate and identify the exact process that causes the functional collapse.

This research paper has a logical structural progression that addresses these concerns: in section 2, we define the geographic location and methodology for conducting marine surveys and establish spatial sampling regimes and statistical analysis methods. Section 3 will present the results using a correlation matrix and an exponential regression model to define habitat loss threshold levels. Section 4 will discuss the mechanisms of coastal habitat loss and assess the role of MPAs in the long-term protection of coastal systems. Finally, section 5 will propose comprehensive conclusions and policy recommendations to protect coastal habitats.

II. Materials and Methods




Figure 1: Human Activities and Mitigation Pathways Affecting Marine Habitat Health

The study used a space-for-time substitution design, employing thirty-eight separate marine sampling sites. Monitoring stations were placed on an existing continuum of human activity, from intensively industrialized harbors and armored tourist beaches to remote marine protected areas. Data were collected from three years of monitoring to account for all seasonal variation (e.g., variation in sea surface temperature, larval settlement periods, storm-influenced terrestrial discharge events).

Figure 1 shows a conceptual schematic of the multiple cascading pathways from land-based urban and industrial activities and from commercial marine activities to physical, chemical, and biological impairment of marine coastal ecosystems. Impacts cascade through systems on multiple spatial and temporal scales. Cascading human impacts on marine habitats. Terrestrial and coastal activities (e.g. Coastal urbanization, coastal land reclamation, industrialization and commercial bottom trawling) can trigger numerous stress pathways into marine environments that impact physical substrates (reduced coral cover, seagrass density, benthic rugosity), cause chemical contamination (increased nutrient load, heavy metals, and Suspended Solids), lead to biological responses (decreased macroinfaunal diversity, loss of reef fish communities and disruption of ecological trophic interactions) and management interventions (Marine Protected Areas, buffer zones and ridge-to-reef management). Arrows signify the causal chain of impacts and resilience measures.

Spatial Coastal Mapping and Anthropogenic Stress Indices

Coastal land cover transformations within the 5km terrestrial zone immediately surrounding each marine sampling station were analyzed based on high-resolution imagery. Four categories of land cover were discriminated with supervised classification: (1) artificial impermeable areas (e.g., Urban areas, industrial facilities, seawalls), (2) aquaculture/agriculture, (3) natural coastal vegetation (mangroves, salt marshes, coastal forests), and (4) undeveloped shoreline. Length of artificial shorelines was calculated as a percentage of armored shoreline with respect to total shoreline length. Pressures of the fishing activity were quantified by ship density based on automatically collected identification system (AIS) tracking information at the corresponding sea quadrants.

Benthic Substrate Assessment and Water Quality Metrics

Benthic surveys were conducted by scuba divers, systematically along 3 parallel fixed-depth, 50 m length line-intercept transects at each station using the traditional chain-and-link method to estimate live coral cover, total macroalgal cover, seagrass shoot density, and structural benthic rugosity. Concomitant with the benthic survey, samples of seawater at 1-2 meters depth were taken and analyzed for Total Suspended Solids (TSS), dissolved inorganic nitrogen (DIN), soluble orthophosphate, and traces of heavy metals (copper, lead, cadmium by inductively coupled plasma mass spectrometry (ICP-MS).

Biological Diversity Evaluation

Visual censuses of macrofaunal assemblages and reef fish populations were conducted at each station along the fixed transects, enumerating fish species, their numbers, and sizes to estimate the total fish biomass. Cryptic benthic macroinvertebrates were collected using a 5cmx5cm scoop and artificial substrate units deployed for 6 months. Based on these metrics, a multi-metric Marine Substrate Health Index (MSHI) was constructed that weighted and combined the measures for live coral cover, benthic rugosity, and macroinfaunal richness (only the more sensitive taxa) in a single composite metric on a scale from 0 to 1.

III. Results

The results of the empirical data showed significant positive and negative correlation coefficients between anthropogenic pressures and the condition of the nearby marine habitat. Coastal vegetated reference sites with healthy mangrove habitat and high coastal physical structure had significant positive benthic structure and biomass metrics, while heavily impacted commercial fishing sites with highly developed coastal zones had significant negative values across all variables of the physical and biological condition of the surrounding marine habitat.

Core Correlations Between Human Stressors and Coastal Health Indices

To examine these associations among anthropogenic activity metrics and indices of coastal ecosystem condition a Pearson correlation matrix was performed using data from all 38 monitoring stations to measure the strength of the relationship between anthropogenic parameters such as percent armoring and commercial trawling, percent industrial runoff and ecosystem variables such as percent cover live coral, percent cover benthic structural complexity, dissolved inorganic nitrogen, and the Marine Substrate Health Index. These correlation coefficients are shown in Table 1. Values which were determined to be statistically significant ($p < 0.01$) are identified with asterisks. High percent armoring and percent industrial runoff showed a high negative correlation with live coral cover, benthic structural complexity, and the condition of the marine substrate; mangrove cover was positively correlated with ecosystem function and nutrient buffering. The correlation between anthropogenic factors and ecosystem variables helps demonstrate the interaction between the variables and the range at which one may not affect the other.

Table 1: Correlations Between Human Activities and Marine Ecosystem Health

Anthropogenic Driver	Live Coral Cover (%)	Benthic Rugosity Score	Dissolved Nitrogen (mg/L)	Marine Health Index (MSHI)
Coastal Armoring (%)	-0.71*	-0.65*	0.58*	-0.76*
Trawling Density (hrs/km ²)	-0.54*	-0.83*	0.22	-0.69*
Industrial Runoff (m ³ /day)	-0.68*	-0.41*	0.82*	-0.72*
Mangrove Cover (%)	0.49*	0.52*	-0.61*	0.68*

Table 1. Pearson correlation coefficients (r) indicating the relationship between human activity indices and marine ecosystem health metrics across thirty-eight sampling nodes (* $p < 0.01$).

Statistical outputs are all clearly significant, suggesting that the density of commercial trawl fishing is very negatively correlated with the aggregate benthic rugosity score ($r = -0.83$)-the severity of the destruction of physical substratum structure. Coastal armoring is very negatively correlated with Marine Substrate Health Index ($r = -0.76$). Pristine mangroves are a great indication of ecosystem services, which are very negatively correlated with DIN ($r = -0.61$), indicating their ability to efficiently "soak up" terrestrial nutrients.

Threshold Response of Marine Substrate Health to Coastal Armoring

The most significant finding from this research was the non-linear threshold response apparent from plotting the marine health indices with increasing intensity of man-made structures. The metrics of the biological communities were already critically stressed before the complete urbanization of the shoreline.

Figure 2 shows that there is consistently high and steady (MSHI >78) environmental quality of the marine substrate condition under relatively low coverage (< 18.2% within an ambient coastal armoring, coastal infrastructure / landward context) by coastal artificial structures, such as breakwaters, seawalls, and urbanized structures. The benthic system at this level of environmental quality is still a complex structure supporting various communities of hard corals, seagrass beds, and associated macro-benthic invertebrate species, maintained fish biomass, and ecological trophic functions. Once the proportion of coastal armoring and urbanization goes above the critical ecological tipping point of 18.2% (Figure 2), MSHI has a steep drop as a sign of a non-linear ecosystem collapse; the complicated structure with high live coral cover and structured benthic environment was destroyed and replaced by a simplified environment of algal turf and soft substrates with dominant influence of some opportunistic species. At the reduced level of environmental quality with the increase in artificial coastal surface structures and the dominance of turf algal community, available niches of habitat structure were seriously decreased and the benthic community structure lost complexity and thus had negative influences on the available space for juvenile reef fish and invertebrates as habitats for shelter and nursery; the interactions among trophic pathways of benthic food

webs and hence sustainable supply of fish biomass significantly decreased with loss of the structured nature and diversity.




Figure 2: Marine Substrate Health Response to Coastal Artificial Surfaces

IV. Discussion

Ecological Consequences of Coastal Armoring and Structural Alterations

The dramatic drop in marine habitat health beyond the 18.2% threshold of coastal armoring reveals the susceptibility of shallow marine systems to structural change at the coast. The destruction of complex, natural intertidal boundaries through the installation of impermeable concrete structures removes valuable ecotones mediating nutrient and sediment flow (Barbier et al., 2011). While natural shoreline components facilitate wave energy dissipation through a combination of vegetated buffer or mobile sediment matrix, seawalls physically reflect wave energy downwards. This change in hydrodynamic regime leads to chronic erosion at the base of concrete seawalls; stripping sandy substrata from shallow marine systems and preventing colonization by juvenile seagrass beds or larval coral structures (Crain et al., 2008).

In addition, human development leads to direct habitat fragmentation at the coast. Complex root networks in mangroves and intertidal channels represent crucial nursery grounds providing shelter for juvenile reef fish and macro-invertebrates, in part by providing protection from large predators, while providing an abundance of food (Halpern et al., 2008). Following the clearance of these valuable, species-rich nursery grounds for the development of beachfront property or commercial harbors, an essential habitat connection between the coastal wetlands and marine coral reef systems is severed, resulting in failed recruitment rates and subsequent declines in adult fish stock throughout marine adjacent zones.

Seafloor Disturbance via Industrial Bottom Trawling

The clear and inverse relationship between commercial bottom trawling and benthic rugosity clearly identifies this as a severe, current physical threat to seafloor ecosystems. Bottom trawling constitutes an intensely managed, ecosystem-wide physical disturbance (Dong et al., 2016). This means steel doors, net frames, weights, and other equipment are being dragged at high force across the seafloor (Kaiser et al., 2006), tearing away slow-growing structural biogenic elements such as gorgonian corals, branching bryozoans, and elaborate sponges that collectively constitute microhabitats and serve as foundation species for numerous macrofaunal taxa (Dunic et al., 2021).

By flattening three-dimensional structural communities onto a featureless plain of sandy substrate and mud, benthic habitat complexity is lost, as the former offers numerous shelters for juvenile fish and specialist invertebrates against predators (Tang et al., 2018). In addition to the physical destruction of structural features, bottom trawling directly suspends enormous volumes of fine, anoxic material into the water column, blocking light from benthic algae and burying suspension feeders such as bivalves and corals with inert sediment (Orth et al., 2006).

Conservation Efficacy of Marine Protected Areas and Coastal Buffers

The clear and positive relationships between intact mangrove cover and downstream indicators ($r = 0.52$ & 0.68 , respectively) reveal that coastal natural features act effectively as buffer zones for coastal ecosystem protection (Abdullah, 2025). Intact mangroves and salt marshes both act as essential sediment sinks and nutrient traps, as complex root structures effectively slow overland runoff and deposit it, while nitrates dissolved within agricultural drainage are removed from water flow before they enter sensitive coral reef systems (Grabowski et al., 2014).

Establishment of protected Marine Protected Area (MPA) zones under full protection is one critically important conservation strategy for coastal habitat management (Valiela et al., 2001; Unsworth et al., 2014). Strictly regulated, no-take fishing zones allow structural habitats to recover and gain back their complexity, while established, fully protected MPAs have the 'spillover effect' that benefits the areas immediately outside of protected waters (Worm et al., 2006; Duarte et al., 2025). However, for effective long-term success, marine protected areas are also in great need of rigid, connected land-use planning (Edgar et al., 2014). Allowing urban development and agriculture to expand relentlessly along the coast while aiming to protect only the underwater portion of the ecosystem integrity results in limited success of the protected area itself. Thus, both coastal land and its adjacent marine habitat must be managed as one unified system.

V. Conclusion

This study emphasizes that human activities pose severe, extensive threats to marine habitats and coastal ecosystems. This study was designed and implemented to provide, for the first time, a clear set of thresholds and tipping points governing marine ecosystem response given a particular human pressure gradient. A threshold value at 18.2% of artificial coasting area was identified that leads to a precipitous decline in benthic habitat health. It is also apparent that the use of commercial bottom trawling causes immense, large-scale physical destruction to structural habitat features within marine ecosystems, greatly flattening these essential areas. Encouragingly, these data unequivocally support the hypothesis that protection of natural coastal habitats such as mangroves and salt marsh ecosystems results in protection of downstream marine ecosystems. Consequently, efforts aimed at improving marine habitat quality must move beyond isolated conservation initiatives towards integrated land use planning for the entire river basin or watershed. It is recommended that further action include: a mandatory network of Marine Protected Areas; regulations on the use of artificial structures along the coast; and restrictions on harmful fishing methods, in order to continue safeguarding global marine ecosystems.

References

- [1] Barbier, E. B., Hacker, S. D., Kennedy, C., Koch, E. W., Stier, A. C., & Silliman, B. R. (2011). The value of estuarine and coastal ecosystem services. *Ecological monographs*, 81(2), 169-193. <https://doi.org/10.1890/10-1510.1>
- [2] Crain, C. M., Kroeker, K., & Halpern, B. S. (2008). Interactive and cumulative effects of multiple human stressors in marine systems. *Ecology letters*, 11(12), 1304-1315. <https://doi.org/10.1111/j.1461-0248.2008.01253.x>
- [3] Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., d'Agrosa, C., ... & Watson, R. (2008). A global map of human impact on marine ecosystems. *science*, 319(5865), 948-952. <https://doi.org/10.1126/science.1149345>
- [4] Dong, Y. W., Huang, X. W., Wang, W., Li, Y., & Wang, J. (2016). The marine 'great wall' of China: local-and broad-scale ecological impacts of coastal infrastructure on intertidal macrobenthic communities. *Diversity and Distributions*, 22(7), 731-744. <https://doi.org/10.1111/ddi.12443>
- [5] Kaiser, M. J., Clarke, K. R., Hinz, H., Austen, M. C., Somerfield, P. J., & Karakassis, I. (2006). Global analysis of response and recovery of benthic biota to fishing. *Marine Ecology Progress Series*, 311, 1-14. <https://doi.org/10.3354/meps311001>
- [6] Dunic, J. C., Brown, C. J., Connolly, R. M., Turschwell, M. P., & Côté, I. M. (2021). Long-term declines and recovery of meadow area across the world's seagrass bioregions. *Global Change Biology*, 27(17), 4096-4109. <https://doi.org/10.1111/gcb.15684>
- [7] Tang, J., Ye, S., Chen, X., Yang, H., Sun, X., Wang, F., ... & Chen, S. (2018). Coastal blue carbon: Concept, study method, and the application to ecological restoration. *Science China Earth Sciences*, 61(6), 637-646. <https://doi.org/10.1007/s11430-017-9181-x>
- [8] Orth, R. J., Carruthers, T. J., Dennison, W. C., Duarte, C. M., Fourqurean, J. W., Heck, K. L., ... & Williams, S. L. (2006). A global crisis for seagrass ecosystems. *Bioscience*, 56(12), 987-996. [https://doi.org/10.1641/0006-3568\(2006\)56\[987:AGCFSE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[987:AGCFSE]2.0.CO;2)
- [9] Abdullah, D. (2025). Environmental Stress–Driven Microbiome Dynamics in Coral Holobionts: Implications for Reef Habitat Degradation and Ecosystem Health Assessment. *Journal of Animal Habitat and Environmental Interactions*, 34-41.
- [10] Grabowski, J. H., Bachman, M., Demarest, C., Eayrs, S., Harris, B. P., Malkoski, V., ... & Stevenson, D. (2014). Assessing the vulnerability of marine benthos to fishing gear impacts. *Reviews in Fisheries Science & Aquaculture*, 22(2), 142-155. <https://doi.org/10.1080/10641262.2013.846292>
- [11] Valiela, I., Bowen, J. L., & York, J. K. (2001). Mangrove Forests: One of the World's Threatened Major Tropical Environments: At least 35% of the area of mangrove forests has been lost in the past two decades, losses that exceed those for tropical rain forests and coral reefs, two other well-known threatened environments. *Bioscience*, 51(10), 807-815. [https://doi.org/10.1641/0006-3568\(2001\)051\[0807:MFOOTW\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0807:MFOOTW]2.0.CO;2)
- [12] Worm, B., Barbier, E. B., Beaumont, N., Duffy, J. E., Folke, C., Halpern, B. S., ... & Watson, R. (2006). Impacts of biodiversity loss on ocean ecosystem services. *science*, 314(5800), 787-790. <https://doi.org/10.1126/science.1132294>
- [13] Unsworth, R. K., van Keulen, M., & Coles, R. G. (2014). Seagrass meadows in a globally changing environment. *Marine pollution bulletin*, 83(2), 383-386. <https://doi.org/10.1016/j.marpolbul.2014.02.026>
- [14] Duarte, C. M., Apostolaki, E. T., Serrano, O., Steckbauer, A., & Unsworth, R. K. (2025). Conserving seagrass ecosystems to meet global biodiversity and climate goals. *Nature Reviews Biodiversity*, 1(3), 150-165. <https://doi.org/10.1038/s44358-025-00028-x>

- [15] Edgar, G. J., Stuart-Smith, R. D., Willis, T. J., Kininmonth, S., Baker, S. C., Banks, S., ... & Thomson, R. J. (2014). Global conservation outcomes depend on marine protected areas with five key features. *Nature*, 506(7487), 216-220. <https://doi.org/10.1038/nature13022>