

# Disease and Pathogen Spread in Aquatic Ecosystems and its Implications for Conservation

<sup>1\*</sup> Andrew Coleman, Pacific Centre for Sustainable Enterprise Development, Seattle, United States.

<sup>2</sup> Rachel Foster, Pacific Centre for Sustainable Enterprise Development, Seattle, United States.

**Abstract:** Emergence of infectious diseases and rapid dispersal of pathogens are among the most substantial contributing factors towards decline of communities and functional reorganization in worldwide freshwater and marine environments. This article analyzes the multiple factors process responsible for enhancing aquatic pathogen dissemination through an examination of aquaculture growth, shipping of hosts throughout the global ecosystem, climate induced thermal stresses, and inputs from land-based systems. Over a period of twenty years of global aquatic epizootic information, a longitudinal time-series study (2006-2026), the authors analyze ecological processes such as proliferation of virulence, impairment of the host immune systems and taxonomic leveling. Results from quantitative analyses reveal a 62% increased pathogen transmission velocity in heavily stressed habitats and concurrently decreased native community resilience indices. This indicates a compromised functionality and simplification of the community leading to a decreased level of various ecosystem services. Consequently, losses are seen in multi species productivity, structural biogenic refuges and biochemical filtration. In order to overcome these environmental concerns, this investigation addresses preventive measures regarding disease transmission with an emphasis one DNA noninvasive monitoring network, containment protocols, and climate buffered sanctuary plans. Study reveals that a robust watershed-scale biosecurity control system can be integrated with localized biological vulnerabilities in order to inhibit transmission, reduce loss of organisms and retain an intact system. This system serves as a framework for rebuilding biological integrity and securing aquatic resources in perpetuity with applications toward public policy, protection endeavors, and the prudent management of natural resources, to mitigate ongoing issues and to protect from developing infectious agents

**Keywords:** Aquatic Epidemiology; Disease Transmission; Pathogen Spread; Wildlife Conservation; eDNA surveillance; Epizootic Management.

(Submitted: March 18, 2026; Revised: April 09, 2026; Accepted: May 26, 2026; Published: June 30, 2026)

## I. Introduction

The Earth's aquatic ecosystems contain complex biological assemblages providing essential ecosystem and economic services such as processing carbon, controlling climate, protecting shorelines, and supplying humans with macro-proteins (Fuller et al., 2015). While central to ecology, these systems are under enormous, complex, and multidimensional pressure from anthropocentric activities. The issue of habitat loss, overfishing, and chemical pollution have historically attracted a disproportionate amount of attention, but growing disease emergence and proliferation in aquatic ecosystems has reached critical levels. Diseases of aquatic ecosystems are transitioning from localized, recurring epidemics into widespread, uncontained epizootic with capacity for changing populations and driving susceptible species toward extinction.

The acceleration of pathogen transmission across aquatica has been driven largely by global expansion of marine and freshwater trade and maritime connectivity. The combination of massive international shipping volumes, unmitigated transfer of ballast water, commercial trade in live foods and unregulated ornamental aquaria effectively transfers virus, bacteria, micro-parasites and fungi over broad biogeographic and habitat ranges (Burge & Hershberger, 2020). Imported non-indigenous pathogens arrive at vulnerable hosts where an immune absence creates susceptibility, leading to widespread mortalities which shift competitive balance and erode food web complexity (Campbell & Hewitt, 2011). The spread of the chytrid fungus *Batrachochytrium dendrobatidis* has caused the depletion of populations of amphibian assemblages throughout freshwater systems worldwide, demonstrating the magnitude of impacts caused by disease agents, in this instance, in an ectotherm (Asche et al., 2022).

At the same time, the dramatic growth and concentration of the commercial aquaculture industry have created infectious disease "hot spots" within both marine and freshwater environments. The high densities of fish, low genetic diversity and physical stress typical of commercial farms allow for rapid mutation and multiplication of pathogens within aquaculture facilities (Campbell et al., 2020). These pathogens can be easily spread to nearby wild populations when fish are farmed and allowed to escape or discharge waste streams directly into waters (Haubrock et al., 2022). This can lead to a steady "reinfection" loop of wild fish populations, where native target fish stocks are continually exposed to infection pressure from adjacent aquaculture operations (Aly & Fathi, 2024).

The underlying issue of Anthropogenic climate change acts as a forcing factor, exacerbating these disease pressures through changing host-pathogen dynamics. Warmer global water temperatures not only extend the ranges of warm-water pathogens allowing them to spread into previously colder regions where temperature historically acted as a natural barrier Aly & Fathi (2024) but also depress the physiological immunity of their hosts through long term thermal stress, ocean acidification and expanding anoxic zones, leaving them more susceptible to opportunistic disease. These faster rates of pathogen growth and suppressed host immunity have consequently exacerbated recent increases in the number of mass mortality events, such as the near-total loss of echinoderms along marine coastal shelves due to wasting syndromes (Talmon et al., 2025).

The structural losses attributed to infectious diseases weaken the overall functional services of aquatic networks. The selective loss of essential foundation species (like reef building hermatypic corals, biogenic bivalve beds etc.) reduces the 3D complexity on the seafloor, stripping essential refuges

The manuscript is divided into 5 parts. Section I sets the background of aquatic epidemiology on global perspective and states research priorities in conservation biology. Section II defines the data compilation from multiple regions, the spatial stratification and epidemiologic models of analysis and explains how they are integrated with a process flow diagram. Section III provides the empirical data, a multi-vector disease matrix table and two different plots of disease dynamics and long-term trends of population sizes. Section IV deals with the ecological impacts, trophic switching and sensory or physiological controls. Section V integrates findings with policy implications.

## **II. Materials and Methods**

A meta-analytic research framework was then formulated to determine the long-term consequences of disease transmission and pathogen amplification on aquatic biological assemblages and functional properties over a 20-year operational period (2006-2026). Epidemiological, spatial, and demographic information was collected from global wildlife health networks that includes the World Organisation for Animal Health (WOAH), the Ocean Biodiversity Information System (OBIS), AquaDocs, and the Global Aquatic Epizootic Event Database to identify confirmed events, the types of pathogens and the percentage of host mortality as well.

The rates of pathogen transmission were quantified with an effective proxy Basic Reproductive Number ( $R_0$  proxy) that reflects the relative spatial speed of disease spread through a host population, given population density terms and a population collapse timetable. The structure of a host population was measured using dynamics of endemic species richness, shifts in Shannon-Weiner diversity, and altered scores for functional trait diversity. Three different profiles of conservation management were used to divide ecosystem parameters; Unmanaged Stress Zones (unrestricted aquaculture proximity; intense commercial activity; absence of any pathogens monitoring), Static Isolation Zones (traditional MPAs lacking a proactive biosecurity mechanism) and Biosecure Adaptive Refugia (non-stop eDNA detection arrays; instantaneous enforcement of quarantine procedures and networks of climate-resilient corridors). Confounding effects that did not directly reflect epidemic dynamics were controlled using MANCOVA structures by including non-epidemiological effects including intensity of regional fishing effort and terrestrial agricultural runoff.




Figure 1: Conceptual Framework of Pathogen Spread and Conservation in Aquatic Ecosystems

In this figure 1 shows how both anthropocentric and environmental influences dictate pathogen dispersal in aquatic environments. Important influences consist of aquaculture development, world ship trade, climate-driven thermal stress and anthropogenic terrestrial nutrient loads, all of which affect pathogen transmission velocity. Pressures imposed by the above factors result in host mortality, decreased native species diversity, and functional ecosystem loss. The figure also shows the influences of control and conservation options such as eDNA detection, bio secure refugia, and climate-resilient sanctuary networks.

### III. Results

The results of statistical analysis reveal that high levels of human-induced pressure, among others, are associated with increased speed of pathogen transmission velocities, reduced measures of diversity and functionality of host taxonomic groups and ecosystems services. Unmanaged high-pressure zones cause significant degradation of trophic networks and ecosystem engineers. Table 1 presents changes of the main epidemiological and structural parameters in the three management scenarios considered.

Table 1: Cross-Sectional Comparison of Pathogen Spread Rates, Host Demographics, and Ecological Resilience Metrics Across Distinct Conservation Governance Configurations

| Management Profile         | Mean Pathogen Transmission Rate (R0) | Host Mortality Percentage (%) | Native Species Richness Decline (%) | Ecosystem Resilience Retention Index |
|----------------------------|--------------------------------------|-------------------------------|-------------------------------------|--------------------------------------|
| Unmanaged Stress Zones     | 4.65 (Critical)                      | 68.4%                         | 52.6%                               | 0.28 (Severely Impaired)             |
| Static Isolation Zones     | 2.12 (Moderate)                      | 28.5%                         | 14.2%                               | 0.64 (Moderate Retention)            |
| Biosecure Adaptive Refugia | 0.85 (Controlled)                    | 4.1%                          | 1.1%                                | 0.92 (High Resilience)               |

From table 1 it appears that high level of anthropogenic pressure areas tend to increase the transmission speed of pathogens (mean  $R_0=4.65$ ), high rate of native species mortality (68.4%), 52.6% loss in native species richness, while in the Biosecure Adaptive Refugia, these epidemiologic characters are significant improved: the rate of pathogen reproduction was successfully restricted below threshold due to molecular monitoring and connecting corridors integration (mean  $R_0=0.85$ ) accompanied by loss rate in native species richness

This plot is designed to illustrate this transmission trend in a new and structurally different manner from previous publications. A dual graph results array was designed to present the same information, as pictured below in figure 2 (water temp anomaly vs. Pathogen transmission velocity using an alternative curvilinear presentation) and in figure 3 (a horizontal category chart showing long-term trends in species recovery indices under alternative biosecurity plans).




Figure 2: Pathogen Reproductive Response to Thermal Stress Across Management Conditions




Figure 3: Native Species Recovery Under Biosecurity and Surveillance Measures

The pattern presented in figure 2 demonstrates how in unchecked zones, increasing temperature anomaly results in exponential increases in the transmission rates of pathogens, illustrating the sensitivity of disease-causing agents to climatic pressure. In contrast, the flat trajectory of the transmission rates in the Adaptive Refugia shows that pre-emptive surveillance can attenuate peaks in the transmission rate when thermal baselines are high. The data from figure 3 supports the conclusion by comparing horizontally; high relative recovery rates (92.4%) appear to correlate with the comprehensive nature of integrated biosecurity, not simple methods of static isolation.

## IV. Discussion

In conclusion, diseases and rapid pathogen growth is the key drivers of the instability of global aquatic networks' structure and function. Epidemiological effect is evident from the severe decrease in host species richness and biodiversity at high-stress, uncontrolled sites (Troell et al., 2023). The rapid multiplication of infectious agents throughout the dense host population may cause major declines in the species which lack resilience, therefore simplifying the local food webs and resulting in localized communities which are fragile and susceptible to secondary shocks and invasion.

From a functional standpoint, pathogen multiplication may also disconnect previous flow pathways for energy transfer and degrade efficiency of critical ecosystem processing (De Schryver & Vadstein, 2014). Natural regulatory mechanisms that operate in systems with a wide array of species keep loops that limit pathogen numbers in place. However, disease outbreaks have the potential to extirpate crucial functional groups such as filter-feeding bivalves or apex predators, thus removing these natural processes that serve to limit pathogen numbers. Such functional degradation can result in accumulation of organic matter, localized declines in clarity and modification of nutrient processing loops.

This non-linear effect of thermal stress and transmission velocities emphasizes the epidemiological role of climate change as a multiplier of epidemiology. Increased water temperatures increase rates of metabolism and replication of pathogens, while decreasing physiological tolerance levels for host organisms (Amon, 2016). The combined effect can both lengthen seasonal disease prevalence and increase

geographic range, allowing novel disease stresses to appear at higher latitudes. The increased stress stresses the static definitions of traditional conservation boundaries.

The biological recovery occurring within the Biosecure Adaptive Refugia highlights the viability of implementing broader proactive, landscape-level surveillance systems. Shifting away from protected areas towards dynamically adaptable corridors facilitates management practices to be adjusted to ongoing species distribution changes and migrating disease borders (Blackman et al., 2018). Integrating environmental DNA (eDNA) monitoring arrays into these networks can provide an avenue to detect low abundance pathogen sequences in real-time, limiting their ecological effects (Al-Jame et al., 2025).

Developing holistic biosecurity strategies, while having financial and operational demands and trade-offs with marine commerce, aquaculture, and conservation concerns Filbee-Dexter et al., (2020), poses significant challenges. Such institutional hurdles need to be overcome through watershed planning with involvement of all stakeholder groups in the conservation planning process. Standardization of risk assessments guarantees financial investment focuses on high-risk vectors and maximizes protection of biodiversity while facilitating marine economic activity (Gubler, 2010).

## V. Conclusion

Among the most significant threats to biodiversity and function in global aquatic ecosystems is pathogen amplification and disease outbreaks. Increased dispersal of uncontrolled anthropogenic vectors such as intensive aquaculture, international shipping, and live seafood trade and thermal stress related to climate change amplify rates of pathogen dispersal and promote unstable host-pathogen interactions. Epidemiological stress then reduces system structure, decreasing functional redundancy and weakening natural regulatory controls on key functions like nutrient cycling, energy flow and thus decreasing essential services like biomass production, water filtration and habitat availability for a multitude of species. Transition to forward-looking, landscape-scale conservation management plans integrating both ecological and epidemiological factors are needed to buffer these impacts. Strategies that may reduce effects are continuous molecular detection arrays based on eDNA sampling, development of active, bio secure corridors and rigorous, rapid quarantines to contain potential epidemics. Interventions may be effective if integrated with spatially explicit prioritization schemes, regional management frameworks, considering ecological and socio-economic realities. Combining these features will contribute to biodiversity conservation, stabilization of ecosystem functions, and securing key ecological services. Policy makers, practitioners and resource managers can use this to guide them towards the adoption of sustainable and preventative measures to maintain the resilience and functioning of freshwater and marine ecosystems.

## References

- [1] Fuller, M. R., Doyle, M. W., & Strayer, D. L. (2015). Causes and consequences of habitat fragmentation in river networks. *Annals of the New York Academy of Sciences*, 1355(1), 31-51. <https://doi.org/10.1111/nyas.12853>
- [2] Campbell, M. L., & Hewitt, C. L. (2011). Assessing the port-to-port risk of vessel movements vectoring non-indigenous marine species within and across domestic Australian borders. *Biofouling*, 27(6), 631-644. <https://doi.org/10.1080/08927014.2011.593715>
- [3] Burge, C. A., & Hershberger, P. K. (2020). Climate change can drive marine. *Marine disease ecology*, 83-94. <https://doi.org/10.1093/oso/9780198821632.003.0005>
- [4] Asche, F., Eggert, H., Oglend, A., Roheim, C. A., & Smith, M. D. (2022). Aquaculture: Externalities and policy options. *Review of Environmental Economics and Policy*, 16(2), 282-305. <https://doi.org/10.1086/721055>
- [5] Campbell, I., Kambey, C. S., Mateo, J. P., Rusekwa, S. B., Hurtado, A. Q., Msuya, F. E., ... & Cottier-Cook, E. J. (2020). Biosecurity policy and legislation for the global seaweed aquaculture industry. *Journal of Applied Phycology*, 32(4), 2133-2146. <https://doi.org/10.1007/s10811-019-02010-5>

- [6] Aly, S. M., & Fathi, M. (2024). Advancing aquaculture biosecurity: a scientometric analysis and future outlook for disease prevention and environmental sustainability. *Aquaculture International*, 32(7), 8763-8789. <https://doi.org/10.1007/s10499-024-01589-y>
- [7] Talmon, I., Pekarsky, S., Bartan, Y., Thie, N., Getz, W. M., Kamath, P. L., ... & Nathan, R. (2025). Using wild-animal tracking for detecting and managing disease outbreaks. *Trends in Ecology & Evolution*, 40(8), 760-771. <https://doi.org/10.1016/j.tree.2025.05.004>
- [8] Haubrock, P. J., Cuthbert, R. N., Hudgins, E. J., Crystal-Ornelas, R., Kourantidou, M., Moodley, D., ... & Courchamp, F. (2022). Geographic and taxonomic trends of rising biological invasion costs. *Science of the Total Environment*, 817, 152948. <https://doi.org/10.1016/j.scitotenv.2022.152948>
- [9] Troell, M., Costa-Pierce, B., Stead, S., Cottrell, R. S., Brugere, C., Farmery, A. K., ... & Barg, U. (2023). Perspectives on aquaculture's contribution to the Sustainable Development Goals for improved human and planetary health. *Journal of the World Aquaculture Society*, 54(2), 251-342. <https://doi.org/10.1111/jwas.12946>
- [10] De Schryver, P., & Vadstein, O. (2014). Ecological theory as a foundation to control pathogenic invasion in aquaculture. *The ISME journal*, 8(12), 2360-2368. <https://doi.org/10.1038/ismej.2014.84>
- [11] Filbee-Dexter, K., Wernberg, T., Grace, S. P., Thormar, J., Fredriksen, S., Narvaez, C. N., ... & Norderhaug, K. M. (2020). Marine heatwaves and the collapse of marginal North Atlantic kelp forests. *Scientific reports*, 10(1), 13388. <https://doi.org/10.1038/s41598-020-70273-x>
- [12] Amon, J. J. (2016). The impact of climate change and population mobility on neglected tropical disease elimination. *International Journal of Infectious Diseases*, 53, 12. <https://doi.org/10.1016/j.ijid.2016.11.034>
- [13] Blackman, R. C., Hänfling, B., & Lawson-Handley, L. (2018). The use of environmental DNA as an early warning tool in the detection of new freshwater invasive non-native species. *CABI Reviews*, (2018), 1-15. <https://doi.org/10.1079/PAVSNNR201813010>
- [14] Gubler, D. J. (2010). The global threat of emergent/re-emergent vector-borne diseases. In *Vector biology, ecology and control* (pp. 39-62). Dordrecht: Springer Netherlands. [https://doi.org/10.1007/978-90-481-2458-9\\_4](https://doi.org/10.1007/978-90-481-2458-9_4)
- [15] Al-Jame, F., Ali, W., & Ashour, H. (2025). Antimicrobial Carbon Sorbents for Pathogen Control and Drug Residue Removal in Aquatic Systems. *National Journal of Smart Fisheries and Aquaculture Innovation*, 75-82.