Aquatic Ecosystems and Environmental Frontiers, Volume 4, Issue 3, September 2026 *ISSN: 3049-186X*
DOI: https://doi.org/10.70102/AEEF/V4I3/AF260308

Biodiversity Assessment and Habitat Restoration Strategies for Enhancing Resilience of Degraded Freshwater Ecosystems

[1*] Ahmed Hassan, Nile Institute for Advanced Development Research, Cairo, Egypt.

[2] Salma Mahmoud, Nile Institute for Advanced Development Research, Cairo, Egypt.

Abstract: Biodiversity loss is increasing in freshwater systems due to habitat modification, pollution, sedimentation, and loss of riparian vegetation that alter ecosystem functions. This research paper brings together biological assessment and habitat diagnosis to develop an approach to identify the restoration actions needed in degraded freshwater ecosystems. The methodological process involves assessing fish, benthic macroinvertebrates, and vegetation communities, along with analyzing habitat structure and water quality across three degradation categories. Environmental science assessments include riparian cover, fine-sediment cover, channelization, dissolved oxygen, turbidity, and nutrients. Numerical examples of comparisons before and after the interventions include fish richness dropping from 18 to 7 species (a 61.1% reduction) and macroinvertebrate richness decreasing from 32 to 12 taxa. Shannon diversity drops from 2.86 to 1.48, which is a 48.3% reduction. Abundance of sensitive macroinvertebrates falls from 42.5% to 9.6%, while that of non-native fish rises from 6.8% to 31.2%. Riparian cover drops from 78% to 19%, fine-sediment cover rises from 15% to 64%, and dissolved oxygen falls from 7.8 to 4.2 mg/L. The above numerical examples illustrate how the framework can be applied, not the actual data collected from the field. The framework links diagnosed pressure to pollution reduction, native vegetation restoration, sedimentation control, and appropriate connectivity interventions. Verification requires seasonal replication and monitoring of restored sites compared with intact sites.

Keywords: Freshwater Biodiversity; Habitat Degradation; Ecological Restoration; Ecosystem Resilience; Macroinvertebrates; Water Quality; Adaptive Management.

(Submitted: May 18, 2026; Revised: June 27, 2026; Accepted: August 11, 2026; Published: September 30, 2026)

I. Freshwater Degradation and the Need for Ecological Recovery

Freshwater ecosystems have a wide range of biological communities and provide many services such as water purification, nutrient cycling, fisheries production, and flood regulation. Rivers, lakes, streams, and wetlands support livelihoods and connect habitats. Pollution, over-abstraction of water, habitat conversion, invasive species, and changes in flow regimes are among the factors that threaten the integrity of these ecosystems. Combining biodiversity conservation, water quality enhancement, habitat conservation, and natural connectivity restoration is necessary to manage these pressures (Tickner et al., 2020). Habitat conservation and restoration of degraded habitats offer an initial step towards safeguarding freshwater biodiversity and ecological restoration (Piczak et al., 2023).

Degradation of habitats impacts aquatic organisms by altering substratum characteristics, vegetative cover, channel morphology, and availability of habitats for feeding and reproduction. These factors influence the distribution of species, composition of the community, and the functioning of aquatic food webs. Restoration needs to focus on habitat maintenance processes and not only physical reconstruction of the habitats. Hydrologic processes, habitat heterogeneity, and reduced environmental stresses are key aspects of aquatic ecosystem restoration (Geist & Hawkins, 2016). Ecological corridors aid restoration by linking fragmented ecosystems, making them an integral part of freshwater ecosystem management (Lin et al., 2022).

Biodiversity evaluation helps detect patterns of degradation and choose appropriate actions. Richness of species, abundance, evenness, and community composition complement each other in describing the state of ecosystems. However, increased richness of species does not reflect resilience, especially in cases

where tolerant and invasive organisms prevail. Therefore, biological indices should be taken into account along with the habitat and environmental conditions. Studies related to the impact of restoration on biodiversity and ecosystem services reveal the necessity to fill the gap in knowledge in order to evaluate different techniques of restoration (Birk et al., 2025). Fish species can also be considered an important aspect since restoration affects the habitat (Huang & Han, 2025).

Although there have been improvements in the implementation of restoration practices, linking ecological assessments to the choice of interventions remains difficult. Biological inventories and habitat assessments may provide distinct characterizations without providing information on what pressures limit the recovery process and on what interventions should be prioritized. Restoration practices therefore need to be evaluated within a larger context that takes into account the health of the river, its biodiversity and the functioning of the ecosystem (Li et al., 2022).

This study aims to fill the gap in knowledge by combining the assessment of biodiversity, the habitat analysis and the restoration planning of degraded freshwater ecosystems. Specifically, the goals of this study are to characterize aquatic biodiversity, assess the condition of the habitats, explore the relationship between environmental stresses and biological communities, and identify the restoration approaches appropriate for the main causes of degradation. The hypothesis of this study is that the improvement of habitat diversity, connectivity, and quality will create the conditions conducive to the recovery of biodiversity.

Section 1 discusses the problem of degradation of freshwater ecosystems and the need for ecological restoration. Section 2 gives an overview of biodiversity and resilience of ecosystems. Section 3 highlights the methods of assessment and restoration planning. Section 4 covers the patterns of biodiversity and degradation of habitats. Section 5 discusses the priorities for restoration and recovery.

Scientific Foundations of Biodiversity and Ecosystem Resilience

Freshwater restoration acknowledges the dynamic nature of rivers and the impacts of hydrology, geomorphology, land use, and climate change on them. It is therefore important to have a flexible design that accommodates changes in the environment. Research on the resiliency of river restoration incorporates practitioner views as well as the assessment of the design. The research highlights the difficulties involved in designing restoration for systems that will continue to change in the future (Tullos et al., 2021).

It is equally important to resolve uncertainties surrounding ecological responses and the effectiveness of management actions for biodiversity conservation. The twenty-five key questions identified in freshwater research address knowledge gaps that limit conservation and restoration activities (Harper et al., 2021). These questions highlight the need for incorporating scientific assessment in the management process. For this case study, it will be necessary to incorporate biodiversity indicators in order to identify ecological constraints and intervention points, acknowledging the fact that one assessment is not enough to measure resilience and recovery after disturbance.

Catchment vegetation serves as another basis for freshwater restoration efforts. Literature on forest restoration indicates that natural regeneration, reforestation, and enrichment planting are some of the methods used to increase biodiversity, reduce soil erosion, and enhance ecosystem services (Sharma et al., 2022). Likewise, literature on native tree restoration in Rwanda addresses issues related to biodiversity enhancement, water resources, and climate change adaptation (Areo et al., 2025). Though these two bodies of literature mostly address the issue of terrestrial restoration, they may be useful in considering the issue of catchment and riparian vegetation in freshwater planning.

Loss of habitat, fragmentation, and degradation form interrelated problems with varied ecological consequences depending on the landscape characteristics. Literature on restoration makes it clear that alterations in biodiversity, functions, and services do not progress in the same way. Functional redundancy may play a part in increasing the resilience of ecosystems; therefore, counting species alone does not help

one to understand ecosystem health (Banks-Leite et al., 2020). The literature on freshwater threats places the issue of biodiversity decline in the context of both local and global factors (Cazzolla Gatti, 2016).

In combination, these experiments have justified the need for combining biological evaluation, habitat analysis, and restoration design. Nonetheless, the different ecosystems where these experiments were conducted, and the various objectives they serve, should be considered when choosing metrics for a given freshwater habitat. The current approach links environmental stressors to relevant strategies, as well as the metrics that can be used to measure them. This approach looks at resilience not as an output of habitat restoration, but as something that needs to be assessed periodically.

II. Ecological Assessment and Restoration Planning Framework

Research Design and Site Classification

The research methodology involves the integration of biodiversity assessments, habitat evaluations, water quality testing, and the selection of priority sites for restoration. Sampling locations are characterized by different levels of freshwater degradation such that biological communities and environmental conditions can be compared along a gradient of disturbance. The site classification includes land use within the vicinity of the site, point sources of pollution, riparian vegetation, channel alteration, and presence of barriers to aquatic connectivity. Locations characterized by low levels of disturbance serve as references, provided that a comparable environment is available.




Figure 1: Methodology Pipeline for Biodiversity Assessment and Freshwater Restoration Planning

The process of site selection, seasonal sampling, biological assessment, and environmental measures is illustrated in figure 1. This method analyzes stressors of degradation and helps in prioritizing the restoration efforts. Monitoring following this analysis assesses the impact on the ecosystem and allows for adaptive management.

Aquatic Biodiversity Characterization

Assessment of biodiversity involves fish, benthic macroinvertebrates, and aquatic vegetation, since these organisms comprise complementary elements of freshwater ecosystems. The fish assessment is conducted through methodologies depending on the depth of the water body and flow rate and any permission for sampling at particular locations. The macroinvertebrates are sampled in a variety of substrates and habitats with either a consistent sampled area or duration of sampling.

The identification of these organisms is made to the lowest possible taxonomic classification levels, with standardization of identification. Species richness, Shannon diversity, and evenness provide information on the diversity of the community, while abundance and taxonomic composition provide information on the changes that cannot be observed in the index of aggregation. The fish abundance is standardized by sampling effort.

Shannon diversity index

$$H' = -\sum_{i=1}^{S} p_i \ln(p_i)$$

Here, H' represents species diversity, S is the number of species, and $p_i = n_i/N$ is the proportion of individuals belonging to species i.

Pielou's evenness index

$$J' = \frac{H'}{\ln(S)}$$

In this case, J' is the index for species abundance evenness and varies between 0 and 1, where high values represent even abundance. The index is used where $S > 1$.

Habitat Condition and Environmental Measurements

Habitat characterization includes substrate composition, sedimentation, bank stability, riparian cover, habitat diversity, and connectivity. Channel width, water depth, and flow velocity provide information on physical parameters influencing habitat availability. Photographs and geographic coordinates serve to mark sampling sites and enable future characterization.

Water quality measurements encompass temperature, pH, dissolved oxygen, electrical conductivity, and turbidity. Nutrient analysis in the laboratory is done following proper sample preservation and analysis methods. Measurements and biological sampling occur simultaneously where feasible. Data quality control is achieved through instrument calibration, replicate samples, and laboratory blanks. All individual habitat parameters are independently meaningful, and any index needs clear scoring criteria and justification before analysis.

Biodiversity and Habitat Relationship Analysis

Descriptive statistics provide summaries of biological and environmental variation by site and season. The differences between degradation classes are analyzed using statistical methods based on the distribution assumptions and the design of sampling. The models include repeated measures of the same site if such data are collected at different seasons.

Community composition is analyzed using ordination with an ecological measure of dissimilarity chosen appropriately. The relationship between biodiversity and environmental variables is analyzed using regression analysis or constrained ordination, after testing for collinearity between predictor variables. Effect sizes and confidence intervals accompany hypothesis testing. Relationships can be used to define the potential ecological constraints but not to prove the cause-and-effect relationship.




Figure 2: Conceptual Model of Freshwater Ecosystem Restoration

Figure 2 highlights the relationship between biodiversity and habitat assessment as well as restoration planning. The proposed actions should help in improving the state of the habitat and recovering the biodiversity. Arrows indicate proposed relationships, while monitoring will be required to see if there are any ecological benefits from restoration.

Restoration Priority Development

Restoration actions are chosen based on the pressures revealed by the assessment process. The lack of riparian vegetation makes it relevant to consider rehabilitation of native vegetation, whereas the high level of sediments indicates the need to find erosion sources and manage the sediments. Barriers to connectivity call for analysis of the restoration opportunities.

The order of intervention takes into account ecological need, benefits to be gained, feasibility, and possible negative consequences. The criteria and weights used for scoring are presented clearly, as well as their sensitivity with respect to different weight alternatives. The proposed benefits are considered the management expectations until verified by monitoring.

Recovery Monitoring and Resilience Evaluation

The restoration assessment method involves taking repeated biological, habitat, and water quality measures at the intervention and control areas. With the use of the method, when there is implementation, the before-and-after approach using sites that have not been subjected to intervention can be used to differentiate the effects of the interventions from broader environmental changes. The factors measured in the process of the evaluation include the native community structure, indicator taxa, habitat connectivity, and seasonality. The resilience assessment approach requires the indication of resilience.

III. Biodiversity Patterns and Habitat Degradation Profiles

Species Richness and Community Diversity

Table 1 illustrates biodiversity indicators in the form of three degradation categories. Biodiversity in the category of fish is 18 and 7 for low and high degradation levels, respectively. This implies that there is a 61.1% reduction. Macroinvertebrates decline in numbers from 32 to 12, while aquatic vegetation declines from 14 to 5. This is what the numerical illustration indicates.

Table 1: Biodiversity Indicators Across Freshwater Degradation Classes

Indicator	Low degradation	Moderate degradation	High degradation
Fish species richness	18	12	7
Macroinvertebrate taxon richness	32	22	12
Aquatic vegetation species richness	14	9	5
Macroinvertebrate Shannon diversity	2.86	2.21	1.48
Macroinvertebrate Pielou's evenness	0.83	0.71	0.60
Sensitive macroinvertebrate individuals (%)	42.5	24.8	9.6
Non-native fish abundance (%)	6.8	17.5	31.2

In table 1, the parameters of richness, diversity, evenness, and community structure are combined. Richness is associated with lower diversity and evenness values, depending on degradation levels. The sensitive macroinvertebrate organisms become rarer in number, while non-native fishes become more abundant. All of the parameters together represent complementary information for describing ecological condition.

Community Composition and Ecological Sensitivity

The Shannon diversity of macroinvertebrates declines from 2.86 to 1.48 on the provided gradient, which is equivalent to a 48.3% decrease. Pielou's evenness decreases from 0.83 to 0.60, suggesting that the number of individuals in each taxon becomes less uniform. Evenness values correlate with the provided richness and Shannon diversity values when rounded.

The number of sensitive macroinvertebrates decreases from 42.5% to 9.6% of macroinvertebrate abundance, which constitutes a decline of 32.9 percentage points. The abundance of non-native fish increases from 6.8% to 31.2% of fish abundance, as shown in figure 3. Such indicators allow differentiation between changes in composition and total abundance. Nevertheless, taxonomic abundance data and proper classification of sensitivity are required for their validation.




Figure 3: Biodiversity Patterns Across Degradation Classes (a) Shannon Diversity and Pielou's Evenness (b) Sensitive Macroinvertebrate and Non-native Fish Relative Abundance

Figure 3(a) illustrates the decrease in Shannon diversity and Pielou's evenness in relation to low to high degradation levels. 3(b) demonstrates the decrease in the relative abundance of sensitive macroinvertebrates and an increase in the abundance of non-native fish. The values provided serve as an example and are not real measurements.

Habitat Conditions and Water Quality

Table 2 presents the habitat and water quality parameters in uniform units. Vegetation coverage within the riparian zone declines from 78.0% to 19.0%, whereas fine sediment coverage rises from 15.0% to 64.0%. Channel modification rises from 8.0% to 71.0% of the channel length that was examined.

Table 2: Habitat and Water Quality Indicators Across Freshwater Degradation Classes

Indicator	Low degradation	Moderate degradation	High degradation
Riparian vegetation cover (%)	78.0	46.0	19.0
Fine-sediment cover (%)	15.0	37.0	64.0
Channel modification (% of assessed length)	8.0	33.0	71.0
Dissolved oxygen (mg/L)	7.8	6.1	4.2
Turbidity (NTU)	8.5	24.7	58.3
Total nitrogen (mg/L)	0.65	1.42	2.85
Total phosphorus (mg/L)	0.03	0.09	0.21

Habitat Structure and Water Quality, along with Biodiversity, are provided in table 2. With more degradation, less cover in terms of vegetation, lower levels of dissolved oxygen, and higher sediment cover, turbidity, and nutrient levels. This way, reporting each indicator individually ensures its ecological significance and assists in recognizing the specific environmental situation that needs further research.

Dissolved oxygen levels vary from 7.8 to 4.2 mg/L, and turbidity varies from 8.5 to 58.3 NTU. Total nitrogen varies from 0.65 to 2.85 mg/L and total phosphorus varies from 0.03 to 0.21 mg/L. Interpretation of actual data depends on temperature, sampling period, freshwater type, and ecological parameters.

Integrated Biodiversity and Habitat Patterns

In the dataset provided, low richness and diversity also correlate with low riparian cover, more sediment deposition, and low dissolved oxygen. Such correlations between biotic and abiotic parameters provide an example of how these variables can be used together. However, they do not indicate which of the pressures is the main one contributing to community dynamics.

Repeating measurements at individual sites are necessary to quantify the relationships and to take into consideration the differences related to seasonality and other uncertainties. Thus, no correlation coefficient or significance value is calculated for these parameters. The connectivity of the habitat must be assessed directly by considering barriers to passage.

Implications for Restoration Planning

The assessment framework is based on the diagnosed pressures and interventions for these pressures. Evidence of riparian vegetation loss leads to evaluation of native vegetation restoration, and excess sedimentation calls for an evaluation of the sources of erosion. Nutrient enrichment points to pollution prevention measures, whereas habitat modification calls for evaluation of habitat heterogeneity and hydrology.

Interventions are determined based on diagnosed conditions, feasibility of ecological restoration, and pressures in the catchment. Baseline assessment determines interventions, and monitoring is used to evaluate biodiversity restoration. Greater resilience needs further evidence of functionality of the restored community.

IV. Restoration Priorities and Pathways to Ecosystem Recovery

Pressure-Based Intervention Selection

Restoration planning identifies degradation stresses and the measures to be taken that target the causes of those stresses. When riparian plant cover is found to be degraded through field survey, the restoration of native vegetation takes precedence. The planting program takes into account the local plant species, bank stability, the need for shade, and maintenance. Sediment deposition calls for identifying the source of erosion and controlling the influx of sediments before habitat restoration.

Habitat Connectivity and Recovery Opportunities

The process of improving connectivity involves determining the type and extent of barriers, the flow conditions, and access to appropriate habitat. Improvements for passage or barrier removal are recommended only where there is more benefit than risk, including that of spreading invasive species. The improvement of habitat involves restoring substrate types, refuge sites, and water flows. The decision of which intervention to choose involves catchment conditions as well.

Adaptive Monitoring and Resilience Evaluation

Rehabilitation priorities take into account ecological necessity, practicality, anticipated benefit, and maintenance considerations. The monitoring process involves comparisons between the biological diversity, habitat condition, and water quality pre- and post-rehabilitation, ideally in the presence of control areas which are not subjected to treatment. Indicators of success include the presence of native species, sensitive species, and long-term community persistence.

V. Conclusion

Degradation of freshwater ecosystems poses risks to biodiversity through the modification of habitats, destruction of vegetation, sedimentation, and declining water quality. The present paper seeks to bridge the gap between assessing the impacts on biodiversity and diagnosing the ecological conditions and restoration options. The illustrative data show large variations among the levels of degradation. The richness of fishes declines from 18 to 7 species, which means a decrease of 61.1%. At the same time, the number of macroinvertebrates drops from 32 to 12 taxa. The Shannon diversity drops from 2.86 to 1.48, a decrease of 48.3%. The number of sensitive macroinvertebrates drops from 42.5% to 9.6%, and non-native fishes increase from 6.8% to 31.2%. Moreover, the percentage of riparian cover declines from 78% to 19%, the fine-sediment cover increases from 15% to 64%, and the dissolved oxygen decreases from 7.8 to 4.2 mg/L. The most important point to remember is that the restoration strategies should take into account the forces causing degradation by reducing pollution, planting native plants, controlling the amount of sediment, and making proper connections. Biodiversity measures serve as a complement to environmental measures, enabling open decision-making on management. Seasonal repetitions, field experiments, and comparison with unmanipulated areas are required to ensure recovery.

References

[1] Tickner, D., Opperman, J. J., Abell, R., Acreman, M., Arthington, A. H., Bunn, S. E., ... & Young, L. (2020). Bending the curve of global freshwater biodiversity loss: an emergency recovery plan. *BioScience*, *70*(4), 330-342. https://doi.org/10.1093/biosci/biaa002

[2] Piczak, M. L., Perry, D., Cooke, S. J., Harrison, I., Benitez, S., Koning, A., ... & Creed, I. F. (2023). Protecting and restoring habitats to benefit freshwater biodiversity. *Environmental Reviews*, *32*(3), 438-456. https://doi.org/10.1139/er-2023-0034

[3] Geist, J., & Hawkins, S. J. (2016). Habitat recovery and restoration in aquatic ecosystems: current progress and future challenges. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *26*(5), 942-962. https://doi.org/10.1002/aqc.2702

[4] Lin, Q., Song, Y., Zhang, Y., Hao, J. L., & Wu, Z. (2022). Strategies for restoring and managing ecological corridors of freshwater ecosystem. *International journal of environmental research and public health*, *19*(23), 15921. https://doi.org/10.3390/ijerph192315921

[5] Birk, S., Weigelt, C., Borgwardt, F., & Kail, J. (2025). Freshwater restoration effects on biodiversity and ecosystem services: a Delphi survey. *Restoration Ecology*, *33*(7), e70119. https://doi.org/10.1111/rec.70119

[6] Huang, W., & Han, Y. (2025). Habitat degradation and restoration in aquatic ecosystems: implications for fish populations. *International Journal of Aquaculture*, *15*. https://doi.org/10.5376/ija.2025.15.0017

[7] Li, P., Li, D., Sun, X., Chu, Z., Xia, T., & Zheng, B. (2022). Application of ecological restoration technologies for the improvement of biodiversity and ecosystem in the river. *Water*, *14*(9), 1402. https://doi.org/10.3390/w14091402

[8] Tullos, D., Baker, D. W., Crowe Curran, J., Schwar, M., & Schwartz, J. (2021). Enhancing resilience of river restoration design in systems undergoing change. *Journal of Hydraulic Engineering*, *147*(3), 03121001. https://doi.org/10.1061/(ASCE)HY.1943-7900.0001853

[9] Harper, M., Mejbel, H. S., Longert, D., Abell, R., Beard, T. D., Bennett, J. R., ... & Cooke, S. J. (2021). Twenty-five essential research questions to inform the protection and restoration of freshwater biodiversity. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *31*(9), 2632-2653. https://doi.org/10.1002/aqc.3634

[10] Sharma, A., Tracy, J., & Panwar, P. (2022). Ecological restoration of degraded forests for achieving land degradation neutrality. In *Land degradation neutrality: Achieving SDG 15 by Forest management* (pp. 191-204). Singapore: Springer Nature Singapore. https://doi.org/10.1007/978-981-19-5478-8_11

[11] Areo, I. O., Mupenzi, C., & Ediagbonya, T. F. (2025). Reforestation and Ecosystem Restoration by Native Tree Species for Biodiversity Enhancement, Water Resource Improvement, and Climate Resilience across Rwanda. *Journal of Applied Sciences & Environmental Management*, *29*(9), 2873. https://doi.org/10.4314/jasem.v29i9.24

[12] Banks-Leite, C., Ewers, R. M., Folkard-Tapp, H., & Fraser, A. (2020). Countering the effects of habitat loss, fragmentation, and degradation through habitat restoration. *One Earth*, *3*(6), 672-676. https://doi.org/10.1016/j.oneear.2020.11.016

[13] Cazzolla Gatti, R. (2016). Freshwater biodiversity: A review of local and global threats. *International Journal of Environmental Studies*, *73*(6), 887-904. https://doi.org/10.1080/00207233.2016.1204133