Aquatic Ecosystems and Environmental Frontiers, Volume 4, Issue 4, December 2026 *ISSN: 3049-186X*
DOI: https://doi.org/10.70102/AEEF/V4I4/AF260412

Influence of Coastal Sediment Grain Characteristics on the Distribution of Burrowing Marine Invertebrates

[1*] Kittisak Wongchai, Siam Centre for Professional Excellence, Bangkok, Thailand.

[2] Siriporn Saengsri, Siam Centre for Professional Excellence, Bangkok, Thailand.

Abstract: The physical properties of coastal sediments have been found to be very influential in determining the distribution, density, and function of burrowing marine invertebrates. Sediment grain size, sorting, porosity, permeability, and substrate stability affect the capability of animals to dig burrows, access food, maintain adequate physiological status, and minimize environmental stress. Differences in the values of the aforementioned characteristics can produce either favorable or unfavorable conditions for bivalves, polychaetes, and burrowing crabs in intertidal and coastal environments. This work presents a study into the effect that the physical characteristics of coastal sediment grains have on the distribution of burrowing marine invertebrates. It employs a limited secondary review of relevant scientific literature to accomplish this objective. Sediment grain size influences burrowing efficiency and the distribution of organisms; however, sediment stability, hydrodynamics, and burial or contamination may alter these relations. Sediment irrigation, biogeochemical cycling, and the functioning of the ecosystem are the ecological aspects related to burrowing activities considered in the study. Sediment characteristics should be considered when assessing coastal habitats and when implementing coastal habitat restoration efforts.

Keywords: Coastal sediments; grain size; marine invertebrates; burrowing behaviour; benthic distribution; sediment characteristics.

(Submitted: August 28, 2026; Revised: September 30, 2026; Accepted: November 13, 2026; Published: December 30, 2026)

I. Introduction

Sediments along the coasts are habitats for various communities of marine invertebrates such as bivalves, polychaetes, and burrowing crabs. Physical features of the substrate determine the ability of such organisms to survive, move, feed, and reproduce in these sedimentary habitats. Grain size is one of the physical features of coastal sediments that plays a critical role in determining the compactness, porosity, and permeability of the sediments, among other physical constraints.

The distribution of burrowing marine invertebrates is affected by the interaction of both the properties of the sediment and the environment. Fine sediments have the capability to hold water and organic material, while coarse sediments tend to offer larger pores with different drainage properties. However, neither is preferred since substrate suitability varies among organisms due to their physiological needs and burrowing habits. Different organisms that have varying physical and behavioral traits can thus have varied reactions to similar sediment environments.

Burrowing activities have also played an important role in shaping the physical and biological features of coastal sediments. They affect the texture of the sediments, water movement, and microbial interaction with organic material in the sediments. Hence, the spatial distribution of burrowing activity may help give insight into environmental stress or other information about the environment. This connection between burrowing behavior and sedimentary properties is important for understanding coastal ecology and the geologic record (Dashtgard, 2011).

Whereas past investigations have considered individual species and the nature of sediments, the conclusions are scattered among various taxonomic categories and different areas of investigation. It is necessary to consider all of these factors together to determine how the grain morphology combines with other properties of the sediments and the environment to affect the distribution of the burrowing species.

This research paper is hence a review of scientific literature regarding the effect of physical parameters of coastal sediments on the distribution of burrowing marine invertebrates. The aims of this research are to evaluate the effects of physical parameters on burrowing efficiency, make comparisons of different invertebrate species' responses to different sediments, and determine environmental factors that affect sediment-invertebrate interactions and the ecological importance of such interactions.

The rest of the paper is arranged as follows. In section 2, there is a review of the relationship between sediment properties and burrowing marine invertebrates. In section 3, the method of the secondary research is provided. In section 4, the findings of the thematic review are presented. In section 5, there is an analysis of the importance of these results for ecology. Section 6 provides the conclusions and directions for future research.

II. Literature Review

Sediment Grain Size and Bivalve Burrowing

The size of the sediment grains will affect the physical state that is found in the sediments by burrowing bivalves. This will affect the burrowing capacity of the bivalves since the arrangement of the grains will be a form of resistance to movement through the sediments.

The study done using two different Donax species has shown that sediment grain size plays an important role in burrowing capability, thus the need for the substrate characteristics to match specific behavioural needs of the species (Nel et al., 2001). Another similar study carried out using Donax trunculus species focused on how sediment grain size affected burrowing, growth, and metabolism, which showed that the substrate characteristics affect not only movement but physiological performance as well (De la Huz et al., 2002). These findings establish that sediment suitability should be evaluated through both behavioural and biological responses.

Sediment Characteristics and Polychaete Distribution

Polychaete worms inhabit many different sedimentary substrates along coastlines but employ different modes of burrowing depending on the type of worm and its habitat. The grain size of the substrate may affect the difficulty of penetration and the creation and maintenance of a burrow.

The study of Perinereis aibuhitensis was conducted on the correlation between the grain size of substrates, burrowing and distributional features showing how significant the sediment texture can be for the spatial distribution of this polychaete (Sun et al., 2019). This implies that the compatibility of the sediments depends on the species and grain size should be taken into account along with other factors.

Sediment Characteristics and Burrowing Crabs

Burrowing crabs are an essential part of intertidal coastal ecosystems. The burrows serve as a habitat and shape sediment structure, while distribution of burrows could indicate changes in the properties of the substrate. Burrow formation and stabilization depend on sediment moisture content, composition and stability.

The study carried out at coastal sites regarding the effect of sediments on the density of burrows of ocypodoid crabs shows that the characteristics of sediments are important for the distribution of burrowing crabs (Saher et al., 2018). The density of the burrows can thus be used as an indicator in ecology when taken together with other factors such as those of the sediments.

Sediment Disturbance and Habitat Suitability

Alterations in sediment may influence the suitability of coastal habitats to support burrowing invertebrates. Coastal projects such as beach nourishment and sediment deposition may involve the use of sediments with different grain size characteristics compared to the natural habitat. There have been

instances where fine-grained beach fills have posed restrictions on the burrowing activities of beach invertebrates (Viola et al., 2014).

The impact of sediment burial is also manifested by the change in the substrate depth, oxygen levels, and access to the sediment surface. The reaction of estuarine benthic invertebrates to burial depends on the mobility and adaptability of those (Hinchey et al., 2006). The obtained results imply that sediment disturbance needs to be taken into account along with sediment grain size.

Burrowing and Sediment Ecosystem Functioning

Physical and biological processes by which burrowing animals affect coastal sediment systems include the creation of pathways within sediments for water flow and the modification of the sediment oxygen status and microbial interactions with organic matter (Kristensen & Kostka, 2005).

Hydrodynamic parameters also have an effect on the formation of benthic communities within new sediments of the marine environment. This is because wave action may have a role to play in the stability of the sediments, their re-distribution and colonization process, thereby affecting the conditions for the settlement of invertebrate communities (Hewitt et al., 2003).

Chemical and Structural Influences on Burrowing

The sediment suitability is not solely based on particle size. The chemicals may alter the behavior of the organisms despite no change in the physical characteristics of the sediment. The study on copper contamination in estuarine and coastal invertebrates observed a disturbance in burrowing activity in both a polychaete and bivalve species (Bonnard et al., 2009).

Sediment erodibility is also dependent on the interplay of sediment properties, biological processes, and habitat structures in the vicinity. Differences in these parameters may affect sediment stability and the ease with which they are physically eroded (Joensuu et al., 2018). This clearly indicates the necessity of carrying out an integrated study of sediment texture and other parameters.

III. Research Methodology

Research Design

This research employs a secondary research approach through a selective review and comparison of existing scientific publications. The study focuses on the correlation between the properties of coastal sediments and the patterns of distribution and burrowing behavior of marine invertebrates. There was no field collection of samples, experimental work in laboratories, recruiting of participants, or any biological observations. The results of the research were obtained through an analysis and comparison of the provided scientific publications.

Literature Selection and Analytical Approach

Eleven peer-reviewed scientific papers were analyzed in the review. These scientific works include the studies of bivalves, polychaetes, burrowing crabs, and also other aspects of benthic ecology. The literature was analyzed taking into consideration its relation to the subject such as grain size of sediment, properties of substrate, ability to burrow, distribution of organisms, sediments disturbance, and environmental stresses.

Thematic Classification

Literature was divided into four major analysis dimensions: physical properties of sediments, organism-specific responses for burrowing, environmental changes, and ecological effects. Physical properties consist of grain size, structure of particles, porosity, permeability, and sediment stability. Organism-specific responses comprise the burrowing potential, burrowing density, habitat selection, and physiology. Environmental changes include waves, burial, beach replenishment, and chemical pollution.

The last dimension deals with irrigation of sediments, bio-geochemical processes, and benthic habitat functions.

The methodology of organization and interpretation of literature is illustrated by figure 1.




Figure 1: Methodological Framework for Reviewing the Influence of Coastal Sediment Grain Characteristics on Burrowing Marine Invertebrate Distribution

IV. Results and Thematic Findings

Influence of Grain Size on Burrowing Performance

Sediment grain size is highlighted in the literature review as one of the critical variables affecting the performance of burrowing in coastal marine invertebrates. Grain size and its distribution have an impact on the resistance that the organism will face when burrowing or moving through the sediment. The bivalves and polychaetes can react in different ways to varying grain sizes.

Coarse and fine sediments lead to varied hydrological and mechanical environments. The coarse sediments usually have bigger spaces between their particles, while fine sediments have smaller spaces and higher cohesion. This results in varied drainage, compaction of the sediments, and the energy needed to dig through them. The applicability of the grains will depend on the organisms and the environment.

Differences Among Invertebrate Groups

According to the analyzed research papers, sediment properties affect various taxonomic groups via different biological processes. The bivalves are impacted due to the resistance experienced during burrowing and substrate properties that affect their stability. The polychaetes are impacted due to the substrate properties in relation to their locomotion and burrow construction process. The burrowing crabs are impacted based on the physical properties of the sediments.

These dissimilarities imply that a single grain-size requirement for sediments is inadequate for all burrowing organisms. The reactions of individual species are necessary to understand the distribution patterns of benthic organisms. Evaluations of habitats need to make such a distinction among different taxonomic groups instead of making them equal to each other.

Effects of Sediment Disturbance and Environmental Stress

Sediment disturbance is capable of changing the connection between grain characteristics and the distribution of the organisms. Beach nourishment and sediment deposition result in the presence of materials whose physical qualities are different from the initial ones. It will make the habitat for organisms worse because of reduced burrowing suitability. Also, sediment burial in large amounts can make it hard for organisms to reach the surface.

Environmental stress can also have an impact on burrowing behavior. Chemical pollution may affect the behavior of animals, while wave action will help in the redistribution of the particles. Thus, it becomes clear that the assessment of the grain size of the sediments needs to be done in the context of the environmental stresses.

Ecological Consequences of Burrowing Activity

Burrowing behavior is part of the ecological function of coastal sediments. Burrows enhance water exchange between the sediment and the environment above it, affect oxygen distribution, and establish microhabitats for microbial populations. Such behavior affects organic matter decomposition and nutrient dynamics in benthic ecosystems.

Burrower abundance and their distribution can, as a result, impact the structure and processes within the ecosystem. If there is any decrease in the number of burrowers, then such a decrease may impact the process of irrigation of the sediments and biological alteration of the sediments. Burrow distribution does not only have implications for the existence of the species but also for sedimentary ecosystem functioning.

Summary of Sediment–Organism Relationships

The key relationships found through the literature review are summarized in table 1. The table describes the major sediment properties and their possible impacts on the burrowers and their ecosystem as a whole.

Table 1: Major Relationships Between Coastal Sediment Characteristics and Burrowing Marine Invertebrates

Sediment or environmental factor	**Influence on burrowing organisms**	**Potential ecological significance**
Grain size	Affects penetration, movement and burrow construction	Influences species distribution and habitat suitability
Particle sorting	Modifies substrate uniformity and pore arrangement	Affects sediment structure and physical stability
Porosity and permeability	Influences water movement and sediment conditions	Modifies oxygen exchange and sediment processes
Sediment stability	Determines the persistence of burrows	Influences habitat continuity
Sediment burial	Can restrict movement and access to the surface	May alter benthic community composition
Wave activity	Redistributes particles and modifies substrate conditions	Influences colonization and habitat stability
Chemical contamination	Can interfere with normal burrowing behaviour	Creates additional environmental stress
Biological burrowing activity	Modifies sediment structure and irrigation	Influences biogeochemical interactions

As is shown in table 1, grain size is a significant factor in habitat suitability, but its influence is associated with other physical, chemical, and biological characteristics. Thus, the distribution of the burrowers can be best explained by the analysis of the sediment conditions and the environment.

V. Discussion

From the literature review above, it is evident that there exists a direct relationship between sediment characteristics in terms of grain size and the presence or burrowing behavior of marine invertebrates. The grain size determines the mechanical characteristics of the sediments, such as packing and resistance to penetration. All these can influence the ability of an organism to be able to dig its burrow or not. The reactions of bivalves, polychaetes, and burrowing crabs vary depending on the biological adaptations.

These results also reveal that sediment grain size alone is insufficient to account for the distribution of burrowing organisms since porosity, permeability, sediment stability, and hydrodynamics all affect the physical environment in which these organisms reside. Burial by sediments, beach replenishment, and chemical pollution are other elements that affect the habitat quality of organisms. It is possible that these environmental factors interact to create varied biological reactions despite the similar grain sizes in the sediments.

There are burrowing animals that have effects on the sediment environment where they live. Burrowing helps in promoting sediment and water exchanges. This interaction shows that the sediment conditions influence the organisms and the organisms influence the conditions of the sediment at the same time. The importance of this interaction lies in the fact that it applies to the coastal ecosystem.

In terms of management, the placement or redistribution of sediments through coastal engineering processes should be assessed based on biological acceptability. Beach nourishment and sediment substitution projects should look into the compatibility of the new sediments based on the ability of the organisms to burrow. In monitoring projects, the integration of burrow densities, species distribution, and sediment characteristics will offer a better indicator of benthic habitat condition compared to just using the size of the sediment grains alone.

There are some limitations in this review. First of all, the review is based on the analysis of a limited number of articles provided by the researcher. Second of all, the reviewed papers were conducted in relation to different species, different geographical locations, and different experimental conditions, which means that there can be no comparison between all results of the reviewed articles. There is no quantitative meta-analysis conducted as well since there is no information about any comparable measurements.

VI. Conclusion

Coastal sediment grain properties can affect the occurrence and burrowing capability of marine invertebrates. Grain size affects the resistance of the substrate, grain structure, and the physical parameters necessary for the formation of burrows. Studies on the topic show that bivalves, polychaetes, and burrowing crabs have different responses depending on their biological and environmental adaptations. Other factors include stability of the sediments, porosity, permeability, wave action, burial and chemical pollution.

Importance of Sediment-Organism Interactions in EcologyThe ecological importance of sediment-organism interactions is not restricted to the spatial distribution of individual organisms. Their burrowing behavior affects the irrigation of sediments, their physical structure, and biogeochemical processes involved. Thus, coastal benthic ecosystem functions depend on interactions between organisms and sediments. Assessment of coastal habitats should take into account properties of sediments, burrowing needs of individual organisms, and environmental stress. Future research should focus on standardization of sediment analysis methods and comparative analysis in different coastal habitats.

References

[1] Dashtgard, S. E. (2011). Linking invertebrate burrow distributions (neoichnology) to physicochemical stresses on a sandy tidal flat: implications for the rock record. *Sedimentology*, *58*(6), 1303-1325. https://doi.org/10.1111/j.1365-3091.2010.01210.x

[2] Nel, R., McLachlan, A., & Winter, D. P. (2001). The effect of grain size on the burrowing of two Donax species. *Journal of Experimental Marine Biology and Ecology*, *265*(2), 219-238. https://doi.org/10.1016/S0022-0981(01)00335-5

[3] De la Huz, R., Lastra, M., & López, J. (2002). The influence of sediment grain size on burrowing, growth and metabolism of Donax trunculus L. (Bivalvia: Donacidae). *Journal of Sea Research*, *47*(2), 85-95. https://doi.org/10.1016/S1385-1101(02)00108-9

[4] Sun, T., Liu, C. E., Li, X., An, D., Yu, H., Ma, Z., & Liu, F. (2019). The effect of substrate grain size on burrowing ability and distribution characteristics of Perinereis aibuhitensis. *Acta Oceanologica Sinica*, *38*(12), 52-58. https://doi.org/10.1007/s13131-019-1348-z

[5] Saher, N. U., Qureshi, N. A., & Siddiqui, A. S. (2018). Influence of sediment characteristics on density and distribution of Ocypodoid crab burrows (superfamily: Ocypodoidea) along the coastal areas of Pakistan. *Acta Ecologica Sinica*, *38*(3), 234-241. https://doi.org/10.1016/j.chnaes.2017.08.006

[6] Viola, S. M., Hubbard, D. M., Dugan, J. E., & Schooler, N. K. (2014). Burrowing inhibition by fine textured beach fill: implications for recovery of beach ecosystems. *Estuarine, Coastal and Shelf Science*, *150*, 142-148. https://doi.org/10.1016/j.ecss.2013.09.003

[7] Hinchey, E. K., Schaffner, L. C., Hoar, C. C., Vogt, B. W., & Batte, L. P. (2006). Responses of estuarine benthic invertebrates to sediment burial: the importance of mobility and adaptation. *Hydrobiologia*, *556*(1), 85-98. https://doi.org/10.1007/s10750-005-1029-0

[8] Kristensen, E., & Kostka, J. E. (2005). Macrofaunal burrows and irrigation in marine sediment: microbiological and biogeochemical interactions. *Interactions between macro-and microorganisms in marine sediments*, *60*, 125-157. https://doi.org/10.1029/CE060p0125

[9] Hewitt, J. E., Cummings, V. J., Ellis, J. I., Funnell, G., Norkko, A., Talley, T. S., & Thrush, S. F. (2003). The role of waves in the colonisation of terrestrial sediments deposited in the marine environment. *Journal of Experimental Marine Biology and Ecology*, *290*(1), 19-47. https://doi.org/10.1016/S0022-0981(03)00051-0

[10] Bonnard, M., Romeo, M., & Amiard-Triquet, C. (2009). Effects of copper on the burrowing behavior of estuarine and coastal invertebrates, the polychaete Nereis diversicolor and the bivalve Scrobicularia plana. *Human and Ecological Risk Assessment*, *15*(1), 11-26. https://doi.org/10.1080/10807030802614934

[11] Joensuu, M., Pilditch, C. A., Harris, R., Hietanen, S., Pettersson, H., & Norkko, A. (2018). Sediment properties, biota, and local habitat structure explain variation in the erodibility of coastal sediments. *Limnology and Oceanography*, *63*(1), 173-186. https://doi.org/10.1002/lno.10622