Aquatic Ecosystems and Environmental Frontiers, Volume 4, Issue 4, December 2026 *ISSN: 3049-186X*
DOI: https://doi.org/10.70102/AEEF/V4I4/AF260415

Ecological Interactions Between Mangrove Leaf Litter and Detritus-Feeding Coastal Organisms

[1*] Matteo Ferri, Renaissance Innovation Development Centre, Florence, Italy.

[2] Elena Moretti, Renaissance Innovation Development Centre, Florence, Italy.

Abstract: Mangrove ecosystem is essential in preserving coastal ecological equilibrium by means of nutrient recycling, organic matter decomposition, and creation of habitats. The leaf litter of mangroves is one of the important sources of organic carbon and nutrients required for detritus feeders such as crabs, amphipods, molluscs, polychaetes, and microorganisms. Nevertheless, the quality of leaf litter, rates of decomposition, environmental conditions, and preferences of diet affect the effectiveness of the ecological interaction. The necessity of the present research emerges due to the increase of coastal disturbances and insufficient knowledge about the influence of characteristics of mangrove leaf litter on food web of the detritus. The uniqueness of the present research project is connected with the creation of the ecological model that links leaf litter production, its decomposition, microbial modification, and feeding behavior of detritivores. The methodological approach suggested is based on comparative ecological assessment of litter production, decomposition experiments, taxonomic classification of organisms, their feeding behavior, and physicochemical properties of water and sediments. This review attempts to explore the impact of litter quality and environmental variability on the detritivores' feeding process as well as the movement of organic material among different mangrove sites. It is believed that the suggested framework will help better understand the role of detritivores in nutrient cycling, sedimentation, and food-web dynamics within the mangrove ecosystems.

Keywords: Mangrove Leaf Litter; Detritus-Feeding Organisms; Organic Matter Decomposition; Nutrient Cycling; Coastal Biodiversity; Microbial Conditioning; Benthic Food Webs; Ecosystem Functioning.

(Submitted: August 31, 2026; Revised: October 01, 2026; Accepted: November 18, 2026; Published: December 30, 2026)

I. Introduction

The mangroves are productive ecosystems on the coasts that sustain diversified biological community and play an essential role in shoreline protection, carbon fixation, and nutrient cycling (Thongtham & Kristensen, 2005). Ecological processes occurring in the mangroves include constant production of leaf litter which is an essential part of organic matter input into coastal ecosystems (Kristensen, 2008; Zaman & Jahan, 2013; Yang et al., 2025). The leaf litter of mangroves decomposes physically, biologically, and microbially, providing nutrition for many detritus feeding organisms. The organisms consist of crabs, amphipods, gastropods, polychaetes, and other benthic invertebrates that take part in the organic matter cycling (Feng et al., 2018). Ecological factors that influence interaction of mangrove leaf litter with detritus feeding organisms are varied. It includes chemical composition of the leaves, decomposition stage, salinity, temperature, oxygen concentration, and microbial activities. Organisms present in decomposing leaves contribute to making them nutritionally available by converting the more complex organic molecules into easily assimilable substances. The next step is decomposition by detritivores, which leads to nutrient recycling and transfer of organic matter through the food web to higher trophic levels. While being ecologically important, shifts in the environmental parameters in the coastal region and deterioration of mangrove ecosystems may affect leaf litter decomposition and food web processes involving detritus (Whitfield et al., 2022; Suhana & Hameed, 2024). To comprehend the above, it is necessary to assess productivity of the ecosystem and ensure coastal biodiversity (Harbour et al., 2021). This research suggests an integrated approach to studying leaf litter properties, processes of decomposition, and feeding by detritivores (Boudouresque et al., 2025).

Problem Statement

The leaf litter from mangroves acts as a basic source of organic material for the coastal detritus feeders, but changes in litter composition, conditions for decomposition of the organic material, and feeding behavior of the organisms may affect nutrient cycling and organic matter dynamics. Habitat degradation, pollution, changes in salinity, and different environmental conditions can have further effects on these ecological processes. The existing studies in ecology tend to look at decomposition of the leaf litter and detritivore community in isolation. Thus, there is a need for a comprehensive ecological approach in this context.

Structure of the Investigation

The study involves an analysis of the interaction of mangrove leaf litter with detritivores in the coastal areas using different inter-related analysis components. The literature review gives the existing ecological processes and constraints within research. The methodology section gives the analysis framework of the ecology study, sampling, and evaluation criteria. Main study involves litter breakdown and organism interaction. Lastly, the discussion, recommendations, challenges, and conclusion sections give the ecological significance and conservation.

II. Literature Review

Mangrove leaf litter decomposition is a vital ecological process for organic matter transformation and nutrient recycling in tropical and subtropical coastal ecosystems. The process of leaf shedding adds significant amounts of plant material into the sediments of mangroves and their neighboring water bodies. Mangrove leaf litter decomposition usually includes processes such as leaching of solutes, microbial colonization, enzymatic degradation, and physical degradation by detritivores. This determines the nutrient content and availability of the litter in the coastal ecosystem. Detritivores are involved in mangrove leaf litter decomposition by means of consumption, fragmentation, mixing of sediments, and redistribution of nutrients. In the process of leaf litter decomposition, crabs tend to fragment leaves and carry organic matter into their burrows, while amphipods and gastropods consume decaying plant matter and the accompanying microorganisms.

Microbial conditioning is also an important factor involved in litter decomposition. The presence of bacteria and fungi on the leaves results in chemical alteration of the leaves and makes some essential nutrients available for the detritivores. Structural properties of the leaves, the percentage of lignin, nitrogen, and the ratio between carbon and nitrogen play important roles in determining decomposition rates and feeding preferences. Environmental factors such as salinity, temperature, tidal flow, and oxygenation of the sediments also play an important role in regulating these factors. Ecological studies conducted in the past have already established the significance of mangrove litter as a source of energy in coastal ecosystems and benthic diversity (Selvaraj, 2000).

Unresolved Areas in Existing Ecological Research

Though a lot of literature exists about mangrove litter decomposition and detritivores feeding, there has been a lack of integration between both these processes. Most studies focus only on litter decomposition rate, microbial colonization or the number of organisms but do not integrate all these processes with respect to nutrient turnover. There is also very little comparative data on species-specific quality of litter, feeding habits of detritivores and the influence of the environment. The interaction between fragmentation of organic matter by organisms and nutrients is yet another area where more ecological evaluation is needed.

III. Methodological Framework

Proposed Ecological Assessment Architecture

In the described approach, the following methods are involved: leaf litter collection, physicochemical analysis, microbial conditioning, and observations of detritivores. At first, the litter samples are gathered at certain mangrove sites and sorted according to the plant type and decomposition level. Lab experiments are conducted to determine the changes in litter weight and its composition; at the same time, the organisms are identified and studied for their behavior while feeding on the litter samples. At all times, the environmental factors are registered.

Research Design and Assessment Procedures

The experimental design is a comparative ecological investigation that uses field studies and controlled laboratory studies. Sites for mangroves will be chosen based on vegetation, tidal action, and sediment features.

It entails:

- Litter collection: Fallen fresh leaves of chosen mangrove species are collected and grouped.
- Decomposition experiments: Litter bags holding the known amounts of leaves are put in representative environments and recovered at regular intervals.
- Identifying the organisms: Detritus-feeding organisms are collected and identified based on their morphological characters.
- Diet study: Controlled study identifies preferences of organisms in litter types and stages of decomposition.
- Environmental measurements: Salinity, temperature, pH, dissolved oxygen, and sediment measurements are made.
- Analysis of data: Statistical comparison and correlation analyses will help understand relations between environmental factors, litter decomposition, and activities of the detritivores.

Proposed Ecological Workflow




Figure 1: Integrated Workflow for Assessing Mangrove Leaf Litter and Detritivore Interactions

Figure 1 demonstrates the sequence of stages in the study of ecology, starting from the selection of mangrove habitats and litter collection. Chemical analysis and decomposition tests will determine the characteristics of litter, whereas microbial colonization and detritivores' feeding will reveal biological processes. Ecological parameters will be defined with the help of measurements. Finally, the statistical analysis of all these processes will demonstrate how those are interrelated.

IV. Ecological Processes and Biological Relationships

Mangrove Leaf Litter Decomposition and Nutritional Transformation

The fallen leaf litter is an important source of organic carbon and nutrients in the coastal ecosystem. Various transformations are applied to the organic materials following their deposition to ascertain their ability to serve as food sources for detritivorous animals. First, leaching of soluble components like sugar, minerals, and other soluble compounds occurs following the interaction between the fallen leaves and tidal waters. In this case, there is loss of litter weight as well as transfer of dissolved organic materials to the water bodies surrounding the leaves. The second transformation involves colonization of the leaves by microorganisms which lead to biological decomposition of the organic materials. Microbes produce extra-cellular enzymes which promote the breaking down of structural compounds like cellulose and other carbohydrates.

Fragmentation of matter promotes decomposition since it provides more surface areas for microorganisms to act on. Fragmentation can be brought about by various activities such as tidal action and biological actions. Mangrove plants vary in leaf thickness, toughness, lignin content, and nutritional value and thus vary in terms of decomposition. Availability of the decomposing litter will thus be influenced by both biological factors and environmental factors. Temperature influences the metabolism of microorganisms, salinity the microorganism populations and organism population, and oxygen determines the path of decomposition.

Detritivore Feeding Interactions and Coastal Nutrient Redistribution

Coastal animals that feed on dead organisms form an integral part of the process by which mangrove litter is converted into usable resources. Various organisms, including crabs, amphipods, gastropods, and polychaetes, interact with the decayed leaves in terms of feeding and fragmenting them, burrowing, and sediment ingestion. The activities determine the flow of organic material among the vegetation, sediment, and other nearby aquatic environments. The crabs found in association with mangroves feed on the fallen leaves and break them down into small fragments. Certain species of the crabs take the leaf debris underground in their burrows to be decomposed under new oxygen and moisture regimes. Gastropods could be feeding on decaying plant debris and associated microbial films, while deposit-feeding polychaetes help in ingesting the organic sediments.

The feeding behavior is affected by the nutritional quality of the leaves, their decompositional state, the chemistry and physical toughness of the leaf material. Microbiological conditioning of the litter can lead to preference of certain organisms since it affects the nutritional value of the leaves. Variations in feeding behavior affect the quantity of litter fragmented and included in sediments. The effect of these relationships is that of nutrients recycling and secondary production facilitation by the detritivores. Their feeding activity ensures the cycling of carbon, nitrogen, and phosphorus through the coastal food chains. However, the extent of this contribution depends on the abundance of the organisms and the conditions prevailing in the ecosystem.

Comparative Ecological Assessment

Table 1 provides a summary of the key biological and environmental components necessary for the functionality of mangrove detrital food webs. It links leaf litter production to microbial conversion and detritivores' feeding behavior. The identified species play various ecological roles in the food web through fragmenting, grazing, consumption, and sediment processing.

Table 1: Major Ecological Components and their Contributions to Mangrove Detrital Food-Web Functioning

Ecological component	Primary process	Ecological contribution
Mangrove leaf litter	Organic matter production	Supplies carbon and nutrients
Bacteria and fungi	Microbial conditioning	Improves detrital nutritional accessibility
Crabs	Leaf fragmentation and consumption	Supports sediment organic matter redistribution
Amphipods	Detrital feeding	Transfers organic energy
Gastropods	Surface grazing	Consumes microbial films and decomposing materials
Polychaetes	Sediment ingestion	Processes sediment-associated organic matter
Environmental factors	Regulation of biological activity	Influences decomposition and organism distribution

Graphical Interpretation of Litter Decomposition




Figure 2: Graphical Interpretation of Litter Decomposition

Figure 2 depicts an imaginary representation of the amount of litter mass that would remain after 12 weeks in three different types of leaves of mangroves. In all cases, there is loss of organic matter gradually, but the manner of decomposition varies, showing how characteristics of leaves affect the persistence of mangrove litter.

V. Discussion

Ecological approach suggested here reveals the connection between decomposition of mangrove leaf litter and detritus-feeding organisms living in the coastal zone. Quality of litter, conditioning by microbes, and environmental factors all affect the processes of transformation and feeding preferences of detritivores. Detritivore feeding is involved into these processes through consuming litter, fragmenting it, and working with sediments, which leads to redistribution of nutrients in the environment. The example of decomposition comparison shows that different kinds of leaves may have different persistence of litter, but still there is a need to prove this empirically. There is an opportunity for research using this methodological approach in order to discover the relationship between litter chemistry, abundance of organisms, and environmental factors.

Conservation and Research Priorities

Further research should be conducted by expanding the comparative studies with respect to different types of mangroves, regions, and seasons in order to identify the variations in decomposition rates of the litter and food consumption habits of the detritivores. The long-term study should explore the impact of variation in salinity levels, temperature levels, coastal pollution, and habitat destruction on the microorganisms and benthic organisms. It would also be important to use molecular technology in order to identify the microorganisms that are responsible for transforming the litter into other forms. Isotope analysis could help track the carbon and nutrients transfer in the detritus food chain.

Research Constraints and Ecological Challenges

Environmental factors which affect mangrove detrital interactions make it difficult to do proper ecological analyses. The changing seasons, the tides, and varying sediment types could lead to different decomposition trends depending on where samples are taken. Proper identification of detritus feeders may need the help of specialists who know taxonomy. Controlled experiments in laboratories may not provide the actual feeding behaviors and environmental interactions of these organisms. In addition, the separation of microbial decomposition and organism-caused fragmentation may be problematic when done naturally. Comparisons may be hindered by lack of data and difficulty in accessing samples.

VI. Conclusion

The goal of this project is to explore the ecology of interactions between mangrove leaf litter and coastal detritivorous species and to explain how both contribute to organic matter decomposition, nutrient recycling, and functioning of coastal ecosystems. The novelty of the research is in its combination of features of leaf litter, microbial conditioning, feeding of the detritivores, and environmental conditions into a single ecological assessment system. The suggested research design consists of collecting litter samples in the field, conducting laboratory decomposition experiments, analyzing microorganisms, identifying organisms, recording feeding behavior, and measuring physical and chemical parameters of the environment. Comparing results on the litter weight loss, composition of nutrients, abundance of organisms, and feeding preferences allows obtaining systematic information about mangrove detrital interactions.

From the concept analysis, one of the key mechanisms involved in the transformation of organic debris from the mangroves includes microbial colonization as well as detritivores. Variations in leaf characteristics and environmental conditions are likely to have some impact on the decomposition processes and the release of nutrition resources. The sample graph below shows the approach through which the comparison of the persistence of leaf litter would be made with respect to various types of mangrove leaves. Nevertheless, the values used in the graph are not factual and no experimental results are presented in the current discussion. In all likelihood, the expected result will be an enhanced understanding of the role played by the detritivores in nutrient cycling and sediment enrichment among others.

References

[1] Kristensen, E. (2008). Mangrove crabs as ecosystem engineers; with emphasis on sediment processes. *Journal of sea Research*, *59*(1-2), 30-43. https://doi.org/10.1016/j.seares.2007.05.004

[2] Harbour, R. P., Smith, C. R., Simon-Nutbrown, C., Cecchetto, M., Young, E., Coral, C., & Sweetman, A. K. (2021). Biodiversity, community structure and ecosystem function on kelp and wood falls in the Norwegian deep sea. *Marine Ecology Progress Series*, *657*, 73-91. https://doi.org/10.3354/meps13541

[3] Zaman, M. B., & Jahan, M. S. (2013). Food and feeding habits of mangrove shellfish, Telescopium telescopium (Linnaeus, 1758) in Bangladesh. *Bangladesh Journal of Zoology*, *41*(2), 233-239. https://doi.org/10.3329/bjz.v41i2.23326

[4] Thongtham, N., & Kristensen, E. (2005). Carbon and nitrogen balance of leaf-eating sesarmid crabs (Neoepisesarma versicolor) offered different food sources. *Estuarine, Coastal and Shelf Science*, *65*(1-2), 213-222. 10.1016/j.ecss.2005.05.014

[5] Yang, Y., Liang, Q., Peng, Y., & Paterson, D. M. (2025). Differential mediation of biogeochemical processes through bioturbation by fiddler and sesarmid mangrove crabs. *Marine Pollution Bulletin*, *211*, 117431. https://doi.org/10.1016/j.marpolbul.2024.117431

[6] Feng, J., Huang, Q., Chen, H., Guo, J., & Lin, G. (2018). Restoration of native mangrove wetlands can reverse diet shifts of benthic macrofauna caused by invasive cordgrass. *Journal of applied ecology*, *55*(2), 905-916. https://doi.org/10.1111/1365-2664.12987

[7] Whitfield, A. K., Able, K. W., Blaber, S. J., Elliott, M., Franco, A., Harrison, T. D., & Houde, E. D. (2022). Feeding ecology and trophic dynamics. *Fish and fisheries in estuaries: A global perspective*, *1*, 255-331. https://doi.org/10.1002/9781119705345.ch5

[8] Suhana, S. N., & Hameed, H. (2024). Distribution and Habitat Preference of Tanaids (Peracarida–Crustacea) in the Kadalundi Estuary on the Southwest Coast of India. *International Journal of Ecology and Environmental Sciences*, *50*(3), 341-348. https://doi.org/10.55863/ijees.2024.0012

[9] Boudouresque, C. F., Astruch, P., Belloni, B., Blanfuné, A., Francesiaz, C., Maury, M., ... & Thibaut, T. (2025). The mediterranean Dune–Beach–Banquette Ecosystem, its pivotal role in Land–Sea coupling and the functioning of coastal Systems, and some related management issues. *Sustainability*, *17*(10), 4556. https://doi.org/10.3390/su17104556

[10] Selvaraj, G. S. D. (2000). Studies on the mangrove ecosystem. *Marine Fisheries Research and Management*, 162-176. http://eprints.cmfri.org.in/id/eprint/4212